

## EFFECTS OF PHOTOPERIODS ON THE GROWTH PERFORMANCE, FEED INTAKE AND FEED EFFICIENCY OF GILTHEAD SEA BREAM (*Sparus aurata*) JUVENILES

El-Sherif, M.S.

Animal Prod. and Fish Reso. Dept., Fac. of Agric., Suez Canal Univ., Ismailia, Egypt

### ABSTRACT

In this study, gilthead sea bream (*Sparus aurata*) juveniles weighting  $16.3 \pm 0.3$  g were reared at a density of 40 fish per 200-L fiberglass tanks for 42 days. The experimental treatments were four photoperiods (6L:6D, 12L:12D, 16L:8D and 24L:0D) with constant light intensity 1500 lx on the water surface. Growth performance, Feed intake, feed conversion efficiency (FCE) and survival rate were investigated. The fish were fed twice daily using a commercial diet containing 46% crude protein at a rate of 2.5 - 3% of body weight/day. The growth measurements were recorded at 15-day intervals during the experimental period. Significantly higher weight gain and specific growth rates were observed in fish exposed to a 24L:0D photoperiod followed by 16L:8D, 6L:6D and 12L:12D photoperiods ( $P < 0.05$ ), respectively. Feed intake was significantly higher in fish exposed to 24L:0D followed by 16L:6D, 6L:8D and 12L:12D photoperiods ( $P < 0.05$ ), respectively. The FCE was significantly higher for fish exposed to 24L:0D and 16L:8D than those exposed to 6L:6D and 12L:12D. The carcass contents of all treatments were unaffected ( $P > 0.05$ ) by varying the photoperiod. These results demonstrated that growth performance, feed utilization and survival rate of gilthead sea bream (*Sparus aurata*) juveniles weighting  $16.3 \pm 0.3$  g can be stimulated significantly by using either a long (16L:8D) or continuous (24L:0D) photoperiod which could be carried out for rearing marine fish juveniles.

**Keywords:** Sea bream (*Sparus aurata*) – photoperiod – growth performance - Feed intake - feed efficiency.

### INTRODUCTION

Determination of the optimal environmental conditions for larvae and juvenile fish is necessary to maximize the production in hatcheries (Hart *et al.*, 1996 and FAO, 2006). One of the most important physical parameters for the growth and survival of fish larvae is photoperiod (Chatain, 1994 and Battaglene, 1995). Many marine fish larvae are visual predators and therefore require light for efficient planktivory. However, optimal photoperiod for larval development, growth and survival may differ, and also change with larval ontogeny (Fielder *et al.*, 2002).

Fish growth is influenced by several biotic and abiotic factors. The photoperiod is classified as a directive factor (Ginés *et al.*, 2004) stimulating the endocrine system; for example, it is known to influence circulating growth hormone (McCormick *et al.*, 1995 and Björnsson, 1997). Photoperiod manipulation has been used successfully to improve the growth of juvenile and larval stages of a number of fish species. A long photoperiod can stimulate growth efficiency in different species (Silva-García, 1996; Boeuf and Le Bail, 1999; Kissil *et al.*, 2001; Ginés *et al.*, 2003; Trippel and Neil, 2003 and Biswas *et al.*, 2006). On the other hand, continuous light can enhance the juvenile growth of Black sea turbot (*Psetta maeotica*) (Turker *et al.*, 2005)

and Tambaqui fish (Aride *et al.*, 2006). However, contradictory findings have been demonstrated in some species. In halibut *Hippoglossus hippoglossus*, Hallaråker *et al.* (1995) found no significant differences between the natural photoperiod and a continuous light regime, whereas Jonassen *et al.* (2000) and Simensen *et al.* (2000) observed that continuous light could be used to improve growth in juvenile halibut when reared from 30 to 170 g. Recently, Biswas *et al.* (2005 and 2006) demonstrated that growth performance can be enhanced in juvenile red sea bream reared from 20 to 100 g body weight. However, there is no information on the effect of photoperiod manipulation on the growth performance of juvenile red sea bream from 1 to 30 g. In order to establish a light regime giving optimal fish growth for a complete production cycle, the effect of photoperiod manipulation should be investigated on different sizes of fish.

Previous experiments on gilthead sea bream *Sparus aurata* L. have demonstrated the positive effects of photoperiod on cultivation of larvae by enhancing prey detection in this visual predator (Flos *et al.*, 2002 and Huidobro and Tejada, 2004). Therefore, the objectives of this study were to investigate the effect of photoperiod on the growth, feed intake, feed conversion efficiency and survival of gilthead sea bream juveniles reared in fiberglass tanks (200-L) for 42 days.

## **MATERIALS AND METHODS**

**Experimental fish:** Gilthead sea bream (*Sparus aurata*) juveniles weighting 12-14 g were obtained from a local commercial farm (Al-Wafaa farm, Ismailia, Egypt) and transported to an indoor system at the same farm. Six hundred fish were randomly distributed into two cylindrical fiberglass tanks with a capacity of 300-L, and acclimated to the experimental conditions (for 2 weeks prior to the start of the study).

**Water quality:** The photoperiod in all tanks was set at 12 h light: 12 h dark (12L:12D) during the first week of acclimation. Each tank was continuously supplied with seawater filtered by 80 µm sand filter at a flow rate of approximately 4 L/ min (Biswas *et al.*, 2006). Throughout the 42-day experimental period, the rearing water in each tank was permanently saturated with oxygen by supplying air continuously through air-stones from an air-blower. The dissolved oxygen and pH were 6.2-7.3 mg/L and 6.5-8, respectively. Average water temperature ranged from 25°C to 27.5°C. Average salinity was 32 ppt.

**Experimental diet:** The fish were fed with a commercial diet of 46% crude protein (according to NRC, 1993), produced by the Sinai shrimps 21 Company, Port Said (Table1). The pellets (3mm) were offered by hand to fish twice daily at 9 am and 6 pm (Eroldogan *et al.*, 2008) at a rate of 2.5 - 3% BW/day (Sahin *et al.*, 1999).

The diet was stored at (4°C) during the experimental duration to avoid the nutrients deterioration.

The ingredients composition and proximate analysis of this feed are provided in Table (1).

**Table (1): Ingredients composition and proximate analysis of the diet offered to gilthead sea bream in this experiment.**

Ingredients	% dry weight
Fish meal <sup>a</sup>	59.9
Soluble fish protein concentrate <sup>b</sup>	1.0
Cod liver oil	5.8
Gelatinized starch <sup>c</sup>	29.8
Vitamin premix <sup>d</sup>	1.0
Mineral premix <sup>e</sup>	1.0
Choline chloride (50%)	0.5
Lignin sulphate	1.0
<b>proximate analysis</b>	
Dry matter	95.5
Crude protein	46
Crude fat	11.7
Crude fiber	4.3
Nitrogen free extract (NFE)	29.6
Crude Ash	8.4

a- Triple Nine, Denmark (CP: 78.6% DM; GL: 9.8% DM).

b- Sopropèche G, France (CP: 72.7% DM; GL: 18.0% DM).

c- C-Gel Instant-12016, Cerestar, Mechelen, Belgium.

d- Vitamins (mg kg<sup>-1</sup> diet): retinol, 18,000 (IU kg<sup>-1</sup> diet); calciferol, 2000 (IU kg<sup>-1</sup> diet); alpha tocopherol, 35; menadion sodium bis., 10; thiamin, 15; riboflavin, 25; Ca pantothenate, 50; nicotinic acid, 200; pyridoxine, 5; folic acid, 10; cyanocobalamin, 0.02; biotin, 1.5; ascorbyl monophosphate, 50; inositol, 400 (Pfizer).

e- Minerals (mg kg<sup>-1</sup> diet): cobalt sulphate, 1.91; copper sulphate, 19.6; iron sulphate, 200; sodium fluoride, 2.21; potassium iodide, 0.78; magnesium oxide, 830; manganese oxide, 26; sodium selenite, 0.66; zinc oxide, 37.5; potassium chloride, 1.15 (g kg<sup>-1</sup> diet); sodium chloride, 0.40 (g kg<sup>-1</sup> diet); dibasic calcium phosphate, 5.9 (g kg<sup>-1</sup> diet) (Pfizer).

**The photoperiod regime:** The fish were exposed to the test photoperiods, where they were acclimated for another week before commencement of the rearing trial. Four different light regimes were established with three replicates for each treatment (light: dark, L:D): (6L: 6D followed by 6L: 6D, 12L: 12D, 16L: 8D and 24L: 0D), using fluorescent lamps. Photoperiods were controlled by a 24-h timer (Multi 9, Merlingerin, Germany) (El-Sayed and Kawann, 2004). Light intensity was kept constant at 1500 lx on the water surface throughout the study. After conditioning for three weeks, the fish were starved for 24 h and the body weight was measured. The stocking density was reduced to 40 fish per tank. The initial mean body weight was approximately 16.3±0.3g and the experimental period lasted 6 weeks (from 20<sup>th</sup> June 2008 to 2<sup>th</sup> August 2008). The growth measurements were recorded at 15-day intervals during the experimental period.

At the end of the experiment, twenty fish in each tank were sampled randomly and frozen at - 20° C for whole body proximate analysis.

**Chemical analyses:**

The diet and the fish in all treatments were analyzed for protein, fat, ash and moisture contents by standard methods (A.O.A.C., 1990).

**Parameters used:**

The parameters used were the weight gain (WG), specific growth rate (SGR), feed conversion efficiency (FCE) and survival rate using the following formulae:

**Weight gain (g)**

$$\text{WG} = W_1 - W_0$$

**Specific growth rate (%/d)**

$$\text{SGR} = 100 \times (\ln W_1 - \ln W_0) / t \quad (\text{According to De- Silva and Anderson, 1995})$$

**Feed conversion efficiency (%)**

$$\text{FCE} = 100 \times [\text{wet weight gain (g)} / \text{dry feed intake (g)}]$$

(According to De- Silva and Anderson, 1995)

**Survival rate (%)**

$$= N_i \times 100 / N_0 \quad (\text{According to Harrell et al., 1990})$$

**Where:**

W<sub>1</sub>= Final wet weight (g)

W<sub>0</sub>= Initial wet weight (g)

T = Time interval in days

N<sub>i</sub> = Number of fish at the end

N<sub>0</sub>= Number of fish initially stocked

**Statistical analysis:**

The data obtained in this study were analyzed by one-way ANOVA procedure of statistical analysis system (SAS, 1988). Means were compared by Duncan's new multiple range test (Zar, 1996).

## **RESULTS AND DISCUSSION**

The growth performance, Feed intake and feed conversion efficiency of gilthead sea bream are presented in Table 2. The final body weights showed significant ( $P < 0.05$ ) differences among the treatments, where the highest values were ( $56.22 \pm 1.8$ ,  $50.40 \pm 2.1$ ,  $46.21 \pm 1.9$  and  $42.66 \pm 1.5$ ) in fish exposed to 24L:0D followed by 16L:8D, 6L:6D and 12L:12D photoperiods, respectively. Similar trends were also observed in total weight gain and SGR among the treatments. The fish reared under a 24L:0D photoperiod showed the highest total weight gain and SGR followed by 16L:8D, 6L:6D and 12L:12D photoperiods (Table 2). In this study, the positive effect of long (16L:8D) and continuous (24L:0D) photoperiods on growth performances of gilthead sea bream is similar to the results observed on the same species, where long and constant photoperiods of both 16 hr of light (16L:8D) and continuous light (24L:0D) have improved growth (Kissil *et al.*, 2001, Ginés *et al.*, 2004 and Biswas *et al.*, 2005). In young specimens (from 25 g to around 200 g) long and constant photoperiods have improved sea bream growth (Silva-Garcia, 1996 and Gines *et al.*, 2004).

From the data in Table (2), feed intake differed significantly among the treatments and the highest value was in fish at the 24L:0D group followed by

16L:8D, 6L:6D and 12L:12D photoperiods . During this experiment, the feed intake was greater under long (16L:8D) and continuous (24L:0D) photoperiods in comparison to those obtained in several fish studies such as red sea bream, *Pagrus major* (Biswas *et al.*, 2005 & 2006); largemouth bass, *Micropterus salmoides* (Petit *et al.*, 2003); Atlantic salmon, *S. salar* (Berrill *et al.*, 2003); halibut, *H. hippoglossus* (Jonassen *et al.*, 2000) and Snapper *Pagrus auratus* (Fielder *et al.*, 2002). The higher feed intake may be due to diurnal fishes are more active under long or continuous photoperiods, and having a greater foraging activity when Feed is delivered, or be related to hormonal stimulation of appetite under a long or continuous photoperiod. For example, growth hormone, known to have a positive effect on appetite, increased with increasing day length during smoltification in salmon (McCormick *et al.*, 1995 and Björnsson, 1997).

**Table (2): Growth performance of juvenile gilthead sea bream *Sparus aurata* exposed to different photoperiods (mean  $\pm$  s.d.)**

Parameters	Photoperiods			
	6L:6D	12L:12D	16L:8D	24L:0D
Initial body weight (g)	16.3 $\pm$ 0.1	16.2 $\pm$ 0.4	16.0 $\pm$ 0.2	16.1 $\pm$ 0.3
Final body weight (g)	46.21 $\pm$ 1.9 <sup>c</sup>	42.66 $\pm$ 1.5 <sup>d</sup>	50.4 $\pm$ 2.1 <sup>b</sup>	56.22 $\pm$ 1.8 <sup>a</sup>
Weight gain (g)	29.9 <sup>c</sup>	26.2 <sup>d</sup>	34.4 <sup>b</sup>	40.1 <sup>a</sup>
SGR (%/d)	2.48 $\pm$ 0.03 <sup>c</sup>	2.29 $\pm$ 0.05 <sup>d</sup>	2.74 $\pm$ 0.02 <sup>b</sup>	2.98 $\pm$ 0.07 <sup>a</sup>
Total feed intake (g/fish)	31.8 <sup>c</sup>	28.6 <sup>d</sup>	36.2 <sup>b</sup>	42.3 <sup>a</sup>
FCE (%)	94.0 $\pm$ 0.8 <sup>b</sup>	91.6 $\pm$ 0.7 <sup>b</sup>	95.0 $\pm$ 0.5 <sup>a</sup>	94.8 $\pm$ 0.4 <sup>a</sup>
Survival rate (%)	97.5 $\pm$ 1.5	95.5 $\pm$ 1.4	100 $\pm$ 0.0	100 $\pm$ 0.0

**Values in a row with different letters are significantly different ( $P < 0.05$ ).**

Survival rate (%) ranged between 100 and 95.5% for all treatments (Table 2). The average data indicated that the highest rate of survival was recorded with 24L : 0D and 16L : 8D photoperiods while the lowest was recorded with 12L :12D. However, there were differences in the rate of survival among treatments.

The data confirm that there was no significant difference in the FCE between fish exposed to 24L:0D and 16L:8D photoperiods (94.9 $\pm$  0.4 and 95.1 $\pm$  0.5). Also, the difference was not significant between fish exposed to 12L : 12D and 6L : 6D photoperiods. But the FCE of 24L: 0D and 16L: 8D was significantly higher than in the 12L: 12D and 6L: 6D photoperiods. The faster growth of red sea bream exposed to a long or continuous photoperiod was accompanied not only by greater feed intake but also by significantly higher FCE ( $P < 0.05$ ) (Biswas *et al.*, 2005). The significantly higher feed intake and FCE in fish exposed to 16L:8D and 24L:0D photoperiods may be because the feeding strategy in fish exposed to these photoperiods reflected most closely the times of maximum appetite (Biswas *et al.*, 2005). The time of feeding has also been reported to affect feed intake and growth performance in sea bream (Ogata *et al.*, 2002; Ginés *et al.*, 2003 and Biswas *et al.*, 2006 ),

goldfish (Trippel and Neil, 2003), rainbow trout (Boujard *et al.*, 1995), Atlantic salmon (Cook *et al.*, 2000) and tilapia (Biswas and Takeuchi, 2003 and EI-Sayed and Kawanna, 2004). Silva-Garcia (1996) reported increased growth in immature gilthead sea bream (25-219 g) under long photoperiods. This author observed similar values between 16L:8D and 24L:0D treatments. Azzaydi *et al.* (1999) also demonstrated that improved growth and feed efficiency are achievable if the quantity of feed supplied is modulated in accordance with the time of maximum appetite. The finding that higher growth accompanied with both higher feed intake and FCE under long and continuous photoperiods parallels the findings in other fish studies, such as red sea bream, *Pagrus major* (Biswas *et al.*, 2005 & 2006); largemouth bass, *M. salmoides* (Petit *et al.*, 2003) and haddock, *Melanogrammus aeglefinus* (Trippel and Neil, 2003). On the other hand, Boeuf and Le Bail (1999) noted that growth might be influenced by light through better feed conversion efficiency and not just stimulated feed intake. However, the finding that higher growth accompanied with both higher feed intake and FCE contrasts the findings by Ogata *et al.* (2002).

Whole body composition data are presented in (Table 3), in general, the values were within the normal ranges of other studies on the same species. Similar results were obtained by Biswas *et al.* (2005). They stated that photoperiod has no significant effects on the protein and lipid contents in the meat of sea bream. In previous photoperiod experiments using gilthead sea bream, Eroldogan *et al.* (2008) did not find variation in body composition.

**Table (3): Chemical composition of sea bream (*Sparus aurata*) exposed to different photoperiod regimes for 42 days (% dry matter basis).**

Photoperiod	Dry matter	Crude protein	Ether extract	Crude ash
6L:6D	29.4	68.02	21.42	9.18
12L:12D	29.8	68.12	21.81	8.72
16L:8D	29.5	69.49	22.37	7.79
24L:0D	29.7	70.03	21.21	7.41

\*Values reported are the mean of four replicates.

## CONCLUSION

These results demonstrated that growth performance, feed utilization and survival rate of gilthead sea bream (*Sparus aurata*) juveniles weighting  $16.3 \pm 0.3$ g can be stimulated significantly by using either a long (16L:8D) or continuous (24L:0D) photoperiod which could be carried out for rearing marine fish juveniles.

## REFERENCES

- A.O.A.C. (1990). Official Methods of Analytical Chemists, Washington, D.C., U.S.A.  
Aride, P.H.R. ; Roubach, R. ; Nozawa, S.R. and Val, A.L. (2006). Tambaqui growth and survival when exposed to different photoperiod. Acta Amazonica, 36 (3): 381-384.

- Azzaydi, M. ; Martínez, F.J. ; Zamora, S. ; F.J. Sánchez-Vázquez and Madrid, J.A. (1999). Effect of meal size modulation on growth performance and feeding rhythms in European sea bass (*Dicentrarchus labrax*, L.). *Aquaculture*, 170: 253–266.
- Battaglene, S.C. ( 1995). Induced ovulation and larval rearing of Australian marine fish. PhD Thesis, University of Tasmania, Launceston, Tasmania.
- Berrill, I.K. ; Porter, M.J.R. ; Smart, A. ; Mitchell, D. and Bromage, N.R. (2003). Photoperiodic effects on precocious maturation, growth and smoltification in Atlantic salmon, *Salmo salar*. *Aquaculture*, 222: 239– Biswas, A.K. and Takeuchi, T. (2003). Effects of photoperiod and feeding interval on Feed intake and growth rate of Nile tilapia *Oreochromis niloticus* L. *Fish. Sci.*, 69: 1010–1016.
- Biswas, A.K. ; Seoka, M. ; Inoue, Y. ; Takii, K. and Kumai, H. (2005). Photoperiod influences the growth, Feed intake, feed efficiency and digestibility of red sea bream (*Pagrus major*). *Aquaculture*, 250: 666– 673.
- Biswas, A.K. ; Seoka, M. ; Tanaka,Y. ; Inoue, Y. ; Takii, K. and Kumai, H. (2006). Effect of photoperiod manipulation on the growth performance and stress response of juvenile red sea bream (*Pagrus major*). *Aquaculture*, 258: 350–356
- Björnsson, B.T. (1997). The biology of salmon growth hormone: from daylight to dominance. *Fish Physiol. Biochem.*, 17: 9–24.
- Boeuf, G. and Le Bail, P.Y. (1999). Does light have an influence on fish growth. *Aquaculture*, 177: 129–152.
- Boujard, T ; Gelineau , A. and Corraze, G. (1995). Time of a single daily meal influences growth performance in rainbow trout (*Oncorhynchus mykiss*). *Aquac. Res.*, 26: 341–349.
- Chatain, B. (1994). Abnormal swimbladder development and lordosis in sea bass (*Dicentrarchus labrax*) and sea bream (*Sparus auratus*). *Aquaculture*, 119: 371–379.
- Cook, J.T. ; McNiven, M.A. ; Richardson, G.F. and Sutterlin, A.M. (2000). Growth rate, body composition and feed digestibility/conversion of growth-enhanced transgenic Atlantic salmon (*Salmo salar*). *Aquaculture*, 188: 15–32.
- De-Silva, S.S. and Anderson, T.V. (1995). Fish Nutrition in Aquaculture. St. Edmundsbury Press, Bury St. Edmunds, Subfolk, Great Britain.
- El-Sayed, A.M. and Kawanna, M. (2004). Effects of photoperiod on the performance of farmed Nile tilapia *Oreochromis niloticus*: I. Growth, feed utilization efficiency and survival of fry and fingerlings. *Aquaculture*, 231: 393-402.
- Eroldogan, O.F. ; Suzer, C. ; Tasbozan, O. and Tabakoglu, S. (2008). The effects of rate – restricted feeding regimes in cycles on digestive enzymes of gilthead sea bream, *Sparus aurata*. *Turk. J. Fish. Aquat. Sci.*, 8: 49-54.
- FAO, (2006). State of world aquaculture. FAO Fisheries Technical Paper. No. 500. FAO, Rome. 134 pp.

- Fielder, D.S. ; Bardsley, W.J. ; Allan, G.L. and Pankhurst, P.M. (2002). Effect of photoperiod on growth and survival of snapper *Pagrus auratus* larvae. *Aquaculture*, 211:135–150.
- Flos, R. ; Reig, L. ; Oca, J. and Ginovart, M. (2002). Influence of marketing and different landbased systems on gilthead sea bream (*Sparus aurata*) quality. *Aquac. Int.*, 10: 189–206.
- Ginés, R. ; Afonso, J.M. ; Argüello, A. ; Zamorano, M.J. and López, J.L. (2003). Growth in adult gilthead sea bream (*Sparus aurata*) as a result of interference in sexual maturation by different photoperiod regimes. *Aquac. Res.*, 34: 73–83.
- Ginés, R. ; Afonso, J.M. ; Argüello, A. ; Zamorano, M.J. and López, J.L. (2004). The effects of long-day photoperiod on growth, body composition and skin colour in immature gilthead sea bream (*Sparus aurata* L.). *Aquaculture Research*, 35:1207-1212.
- Hallaråker, H.; Folkvord, A. and Stefansson, S.O. (1995). Growth of juvenile halibut (*Hippoglossus hippoglossus*) related to temperature, day length and feeding regime, *Neth. J. Sea Res.*, 34: 139–147.
- Harrell, R.M.; Kerby, J.H. and Minton, R.V. (1990). Culture and propagation of striped bass and its hybrids striped bass committee, Southern Division, American Fisheries, Society, Bethesda, Maryland.
- Hart, P.R. ; Hutchinson, W.G. and Purser, G.J. (1996). Effects of photoperiod, temperature and salinity on hatchery-reared larvae of the greenback flounder (*Rhombopteryx tapirina* Günther, 1862). *Aquaculture*, 144: 303–311.
- Huidobro, A. and Tejada, M. (2004). Gilthead seabream (*Sparus aurata*): suitability for freezing and commercial alternatives. *J. Sci. Feed Agric.*, 84: 1405–1413.
- Jonassen, T.M. ; Imsland, A.K. and Stefansson, S.O. (2000). Interaction of temperature and light on growth in Atlantic halibut, *Hippoglossus hippoglossus* L. *Aquac. Res.*, 31: 219–227.
- Kissil, G.W. ; Lupatsch, I. ; Elizur, A. and Zohar, Y. (2001). Long photoperiod delayed spawning and increased somatic growth in gilthead seabream (*Sparus aurata*). *Aquaculture*, 200: 363–379.
- McCormick, S.D. ; Björnsson, B.T. ; Sheridan, M. ; Eilertson, C. ; Carey, J.B. and O'Dea, M. (1995). Increased day length stimulates plasma growth hormone and gill Na<sup>+</sup>, K<sup>+</sup>-ATPase in Atlantic salmon (*Salmo salar*). *J. Comp. Physiol.*, 165B: 245–254.
- NRC (National Research Council), (1993). Nutrient requirements of warm water fishes and shellfishes, revised edition. National Academy Press, Washington, D. C., USA.
- Ogata, H.Y.; Oku, H. and Murai, T. (2002). Growth performance and macronutrient retention of offspring from wild and selected red sea bream (*Pagrus major*). *Aquaculture*, 206: 279–287.
- Petit, G. ; Beauchaud, M. ; Attia, J. and Buisson, B. (2003). Feed intake and growth of largemouth bass (*Micropterus salmoides*) held under alternated light/dark cycle (12L:12D) or exposed to continuous light. *Aquaculture*, 228: 397–401.

- Sahin, T. ; Akbulut, B. and Aksungur, M. (1999). Investigation on weight loss occurring in winter season and growth of gilthead sea bream (*Sparus aurata*) reared in the Black Sea. Turkish J. Marine Sciences, 5(2): 79-86.
- SAS, (1988). SAS users' guide. Statistics. A. A. Ray. Ed. SAS Inst., Inc., Cary, NC., 67 (3).
- Silva-García A.J. (1996). Growth of juvenile gilthead seabream(*Sparus aurata* L.) reared under different photoperiod regimes. The Israeli Journal of Aquaculture-Bamidgeh, 48: 84-93.
- Simensen, L.M. ; Jonassen, T.M. ; Imsland, A.K. and Stefansson, S.O. (2000). Photoperiod regulation of growth of juvenile Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 190: 119-128.
- Trippel, E.A. and Neil, S.R.E. (2003). Effects of photoperiod and light intensity on growth and activity of juvenile haddock (*Melanogrammus aeglefinus*). *Aquaculture*, 217: 633-645.
- Turker, A. ; Yigit, M. and Ergün, S. (2005). Growth and feed utilization in juvenile Black sea turbot (*Psetta maeotica*) under different photoperiod regimes. Turk. J. Vet. Anim. Sci., 29: 1203-1208.
- Zar, J.H. (1996). Biostatistical Analysis. Prentic Hall, Upper Saddle River, NJ, USA.

**تأثير فترات الإضاءة على أداء النمو و الغذاء المأكول و كفاءة تحويل الغذاء للطير اليافع لأسماك الدنيس .**

**محمد سعد الشريف**

**قسم الانتاج الحيواني والثروة السمكية - كلية الزراعة - جامعة قناة السويس - الاسماعيلية - مصر**

في هذه الدراسة - تم رعاية الأطوار اليافعة لأسماك الدنيس بمتوسط وزن  $16.3 \pm 3$  جم في تناكل من الفيبرجلاس (٢٠٠ لتر ماء) بكثافة تخزينية ٤٠ سمكة لكل تناكل لمدة ٤٢ يوم على أربع فترات إضاءة مختلفة (٦ ساعات إضاءة: ٦ ساعات إظلام وتكرر ٦ ساعات إضاءة : ٦ ساعات إظلام، ١٢ ساعه اضاءه: ١٢ ساعه إظلام، ١٦ ساعه اضاءه: ٨ ساعات إظلام و ٢٤ ساعه اضاءه: عدم وجود اظلام) . وكانت كثافة الضوء ثابتة (١٥٠٠ لوكس) على سطح الماء وذلك لتغذير أداء النمو، الغذاء المأكول، كفاءة تحويل الغذاء ومعدل البقاء . تم تغذية الأسماك مرتين يوميا على علبة تجارية تحتوى على ٤٦٪ بروتين وبمعدل ٣-٢.٥٪ من وزن الجسم يوميا . تم تقدير قياسات النمو المختلفة كل ١٥ يوم خلال فترة التجربة . أوضحت النتائج وجود اختلافات معنوية عالية لكل من الوزن المكتسب ومعدل النمو النوعي في الأسماك المعروضة لـ ٢٤ ساعة إضاءة ، ثم ١٦ ساعة إضاءة، ثم ٦ ساعات إضاءة، ثم ١٢ ساعة إضاءة ( $P < 0.05$ ) . لوحظ اختلافات معنوية عالية للغذاء المأكول في الأسماك المعروضة لـ ٢٤ ساعة إضاءة ثم ١٦ ساعة إضاءة يليها ٦ ساعات إضاءة ثم ١٢ ساعة إضاءة ( $P < 0.05$ ) . كانت كفاءة تحويل الغذاء أعلى معنويًا للأسماك المعروضة لـ ٢٤ ساعة ضوء والمعرضة لـ ١٦ ساعة ضوء مقارنة بالأسماك المعروضة لـ ١٢ ساعة ضوء والمعرضة لـ ٦ ساعات ضوء . كانت مكونات الجسم لكل المعاملات لم تتغير أو تتأثر معنويًا بتغيير فترة الإضاءة .

وقد خلصت النتائج إلى أن أفضل أداء لنمو الطير اليافع لأسماك الدنيس يكون تحت نظام ٢٤ ساعة إضاءة: عدم وجود إظلام وأيضًا ١٦ ساعة إضاءة : ٨ ساعات إظلام، وهذا يرجع إلى تحسن في الغذاء المأكول،ارتفاع كفاءة تحويل الغذاء ومعدل البقاء - لهذا يوصى بتطبيق ذلك لرعاية الطير اليافع لأسماك المياه المالحة .