

## **EFFECT OF SOME ANTIOXIADNTS AND METHYL- DONORS ON THE PRODUCTIVE AND REPRODUCTIVE PERFORMANCE OF TURKEY HENS**

**Raya , A. H. M.<sup>1</sup> ; F. S. A. Ismail<sup>1</sup>; M. Y. Mostafa<sup>2</sup> and M. A. L. Gorgy<sup>2</sup>**

**1. Poultry Production Dept., Fac. Agric., Mansoura University.**

**2. Animal Production Res. Inst., Agric. Res. Center, Dokki, Giza, Egypt.**

### **ABSTRACT**

The objective of this study was to investigate the effect of using some antioxidants and methyl-donors on the productive and reproductive performance of turkeys. This study was carried out at Mehalet Mousa Animal Production Research Station, Kafr El-Sheikh, belonging to the Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture, Egypt. A total of 168 Bronze female turkeys, 36 weeks of age were used. The birds were randomly distributed into 8 experimental groups, each treatment consisted of three equal replicates. The birds were individually housed in battery cages. The experimental period lasted 12 weeks (from 36 to 48 weeks of age). The birds were fed a basal yellow corn-soybean meal diet, supplemented with two natural antioxidants, vitamin E (alpha-tocopheryl acetate) at 250 mg/kg diet and vitamin C (L-ascorbic acid) at 200 mg/kg diet in addition to Choline (Choline chloride 60%) at 2.25 g/kg diet one of labile methyl group donors, singly or in combination. The results obtained can be summarized as follows. Dietary supplementation with vitamin E had positive effects on feed conversion, hen-day egg production rate ( $P \leq 0.05$ ), egg fertility and total hatchability and length of oviduct of turkey hens as compared to those of the control hens. There were no significant differences among the different dietary treatments in mortality rate, relative weights of egg albumen, egg yolk, yolk index, shell thickness, yolk color, concentrations of blood haemoglobin and serum total protein, hatchability of fertile eggs or in relative weights of carcass traits of turkey hens. It can be concluded that dietary supplementation with vitamin E at 250 mg/kg diet can induce beneficial effects on productive and reproductive performance of turkey hens.

### **INTRODUCTION**

The generation of free radicals and lipid peroxidation can contribute to the development of different diseases in humans as well as animals, with a decrease in the live performance and product quality in poultry. Antioxidants are involved in the prevention of cellular damage, which is commonly responsible for aging and a variety of diseases. Antioxidants can safely interact with free radicals and terminate the chain reaction before vital molecules are damaged. Although there are several enzyme systems within the body that scavenge free radicals, the most important micronutrients serving as antioxidants are vitamin E and vitamin C.

Vitamin E is a natural fat-soluble antioxidant and a major chain-breaking antioxidant in biological systems in birds (Khan *et al.*, 2011). Poultry cannot synthesize vitamin E, therefore, their requirements of this vitamin must be met from dietary sources (Chan and Decker, 1994). Vitamin E has been reported to be an excellent biological antioxidant that protects cells and tissue from lipoperoxidative damage induced by free radicals (McDowell, 1989).

Vitamin E is also known to be a lipid component of biological membranes (Halliwell and Gutteridge, 1989). Vitamin E serves as a physiological antioxidant through inactivating free radicals, increasing feed intake and improving egg production (Kirunda *et al.*, 2001), and egg quality (Puthongsiripon, 1998). Vitamin E requirements in laying hens and other food producing animals have been established to avoid clinical symptoms of deficiency (NRC, 1994). It has been proposed that the recommended levels of dietary vitamin E should vary depending on the susceptibility to oxidation rather than the amount needed to prevent signs of deficiency and mortality (Wang *et al.*, 1996).

Vitamin C is a water-soluble antioxidant, vitamin that poultry can synthesize it, so they do not require dietary source of vitamin C (NRC, 1994). It has been, however, reported that ascorbic acid synthesis and utilization are inadequate under stress conditions such as low or high environmental temperatures, humidity, high productivity, and parasite infestation (McDowell, 1989). Vitamin C is involved in several biochemical processes and its function is also related to its reversible oxidation and reduction characteristics within the cells (McDowell, 1989). Vitamin C exhibits an antioxidant effect by removing free radicals from the environment (Tanaka *et al.*, 1997) while the effective mechanism of vitamin E in its reactions with active radicals is to break chains, to exert pressure, to renovate, and to increase endogen defence (Azzi and Stocker, 2000).

Choline, is,  $\beta$ -hydroxy ethyl trimethyl ammonium hydroxide (Sheard and Zeisel, 1989). Choline is considered as an essential water soluble nutrient and is usually grouped within the B-complex family. It is chemically unique as it has three methyl groups attached to the structure and takes part in various functions in the body as a methyl group's donor.

Choline has three essential metabolic functions Chan (1984) and Kettunen *et al.* (2001) that cannot be performed by other methyl donors in the system which include:

- 1) Structural component of cell membranes, as a constituent of phospholipids (phosphatidylcholine) and thereby playing an essential role in building and maintenance of cell structure.
- 2) Lipotropic agent in fat metabolism in the liver by utilizing and transporting fat and thereby preventing fatty liver.
- 3) Precursor for acetylcholine synthesis, (a neurotransmitter agent) for nerve impulses. NRC (1994) suggested that the choline requirement of turkeys ranges from 800 to 1600 mg/kg feed. The dietary need for choline is known to be influenced by levels of other nutrients involved in methyl group metabolism as well as type or strain, age and physiological status of the bird. This study aimed to study the effect of using some antioxidants and methyl-donors on the productive and reproductive performance of turkey hens.

## MATERIALS AND METHODS

The experimental work of this study was carried out at Mehalet Mousa Animal Production Research Station, Kafr El-Sheikh, belonging to the Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture, Egypt. This experiment was carried out to study the effect of using some antioxidants and methyl-donors on the productive and reproductive performance of turkey hens. A total of 168 Bronze female turkeys, 36 weeks of age were used. The birds were randomly distributed into 8 experimental groups, each treatment consisted of three equal replicates. The birds were individually housed in battery cages fitted with individual feeders and automatic nipple drinkers. The experimental period lasted 12 weeks (from 36 to 48 weeks of age). Birds were fed *ad-libitum* and fresh water was available all the time, during the experimental period. Photoperiod was 16 hours daily during the experimental period.

**Table 1: Composition and calculated analysis of the basal diet used in the experimental diets:**

Ingredients	%
Yellow corn	70.00
Soybean meal (44%CP)	11.50
Fish meal (65%CP)	10.00
Di calcium phosphate	2.00
Limestone	6.00
Salt ( NaCl)	0.30
L-Lysine	0.15
DL.Methionine	0.05
Total	100
Calculated analysis:	
Crude protein %	17.72
Metabolizable Energy(kcal/kg diet)	2920

**The experimental groups in this experiment were fed one of the following diets :**

- T1: a basal diet (control group) without any supplementation.
- T2: basal diet + (vitamin C) 200 mg/kg diet .
- T3: basal diet + (vitamin E) 250 mg/kg diet.
- T4: basal diet + (Choline) 2.25 g/kg diet.
- T5: basal diet + (vitamin C) 200 mg/kg diet + (vitamin E) 250 mg/kg diet.
- T6: basal diet + (vitamin C) 200 mg/kg diet + (Choline) 2.25 g/kg diet.
- T7: basal diet + (vitamin E) 250 mg/kg diet + (Choline) 2.25 g/kg diet.
- T8: basal diet + (vitamin C) 200 mg/kg diet + (vitamin E) 250 mg/kg diet + (Choline) 2.25 g/kg diet .

Egg weight, egg number and mortality rate were recorded daily and feed intake was calculated weekly. Feed conversion (g feed/g egg mass) was also calculated. Egg quality parameters were measured at peak of egg production (42 weeks of age) during the experiment using 80 eggs (10

eggs/treatment). These parameters involved relative weight of egg components (yolk, albumen and shell), and some exterior and interior parameters of egg quality. Egg shell thickness was measured in mm using a micrometer. Egg shape index was calculated as egg width divided by an egg length. Yolk index was calculated as yolk height divided by yolk diameter. Haugh unit was measured according to the formula presented by Eisen *et al.* (1962). Yolk color was determined by using a Roche yolk color fan. Semen was collected from toms housed in floor pens and fed on the same experimental diets and artificially inseminated to hens. The insemination was made as soon as possible after semen collection. The artificial insemination was done once a week. All hens were inseminated intra-vaginally. Raw semen was diluted in the rate of 1:1 with 0.9 % NaCl. The insemination was performed by inserting 0.1 ml of the diluted semen with one millimeter tuberculin syringe into the vagina of each hen. Fertility and hatchability (four hatches) of eggs were made every 3 weeks of the experimental period. Fertility was calculated as the number of fertile eggs relative to total number of eggs set; while egg hatchability was calculated as the number of healthy hatched chicks relative to total number of eggs and of fertile eggs. In this experiment, blood samples were collected from 48-week-old turkey hens (3 specimens per treatment) by venipuncture of the wing vein. Immediately after collection, blood sample was transferred into two tubes, a heparinized tube and non-heparinized one. Heparinized whole blood samples were used for the determination of hemoglobin concentration according to Pilaski (1972) and haematocrit value, as described by Hunsaker (1969). The separated sera from non-heparinized blood samples were used for the colorimetric estimation of levels of glucose (Trinder, 1969), total protein (Henry *et al.* 1974), albumin (Doumas *et al.* 1977), cholesterol according to Watson (1960) and total lipids according to Zollner and Kirsch (1962), using commercial kits. The concentration of serum globulin was calculated by the difference between serum total protein and albumin.

At the end of experimental period (48 wks old), 3 hens from each treatment were randomly selected and slaughtered for carcass evaluation. Carcass was eviscerated and head and shanks were removed. Each portion was expressed as a percentage of live body weight.

**Statistical analysis:**

Data were statistically analyzed by one-way analysis of variance by General Linear Model (GLM) procedures (SAS, 2003). Significant difference among means of treatments were detected by Duncan Multiple Range Test (Duncan, 1955). The following model was used to study the effect of vitamin C, vitamin E, Choline and their combinations on parameters investigated as follows:  $Y_{ij} = \mu + T_i + e_{ij}$

Where,  $Y_{ij}$  = an observation

$\mu$  = overall mean

$T_i$  = treatments ( $i = 1, 2 \dots$  and 8)

$e_{ij}$  = residual "random error".

## RESULTS AND DISCUSSION

### Productive performance of turkey hens:

Data on the effect of dietary single and combined addition of vitamin C, vitamin E and choline on productive performance, in terms of feed intake, hen-day egg production rate, egg weight and feed conversion ratio, of turkey laying hens from 36 to 48 weeks of age are given in (Table 2).

As presented in Table 2, hens fed diet supplemented with vitamin C plus vitamin E consumed significantly more ( $P \leq 0.05$ ) feed compared with the control group. Feed intakes of the remaining experimental groups were not significantly different from that of the control birds. The highest mean of hen-day egg production rate ( $P \leq 0.05$ ) was achieved by hens fed vitamin E-supplemented diet, followed by that of hens fed diet supplemented with vitamin C plus vitamin E ( $P \leq 0.05$ ). Hen-day egg production rates of the other experimental groups were not significantly different from that of the controls. The best means of egg weight ( $P \leq 0.05$ ) were attained by hens fed diets supplemented with vitamin C plus choline, vitamin C alone, vitamin E plus choline and a mixture of vitamin C, vitamin E and choline in a descending order as compared to that of the control turkey hens. The worst mean ( $P \leq 0.05$ ) of egg weight was recorded for hens fed diet supplemented with choline alone compared with that of the controls. Egg weights of the other experimental groups did not differ from that of the control group. The best means of feed conversion ( $P < 0.05$ ) was attained by hens fed diet supplemented with vitamin E alone compared with that of the control hens.

**Table (2): Effects of dietary single and combined addition of vitamin C , vitamin E and choline on productive performance of turkey hens from 36 to 48 weeks of age.**

Items	Feed intake (g) 36-48 wks	Egg Production (%) 36-48 Wks	Egg weight (g) 36-48 wks	Feed conversion (g feed /g egg) 36-48 wks	Mortality (%) of turkey hens
Control (Basal diet)	128.22 <sup>bc</sup>	51.70 <sup>c</sup>	76.30 <sup>bc</sup>	3.24 <sup>ab</sup>	0.00
Basal diet +Vitamin C	133.33 <sup>b</sup>	54.42 <sup>bc</sup>	78.36 <sup>a</sup>	3.12 <sup>ab</sup>	0.04
Basal diet +Vitamin E	133.00 <sup>b</sup>	62.75 <sup>a</sup>	75.21 <sup>cd</sup>	2.81 <sup>c</sup>	0.00
Basal diet +Choline	126.99 <sup>c</sup>	50.56 <sup>c</sup>	73.83 <sup>d</sup>	3.39 <sup>a</sup>	0.00
Basal diet + Vitamin C + Vitamin E	138.94 <sup>a</sup>	57.31 <sup>b</sup>	77.59 <sup>ab</sup>	3.12 <sup>ab</sup>	0.00
Basal diet + Vitamin C + Choline	126.27 <sup>c</sup>	51.47 <sup>c</sup>	79.00 <sup>a</sup>	3.10 <sup>b</sup>	0.00
Basal diet + Vitamin E + Choline	132.27 <sup>b</sup>	53.51 <sup>bc</sup>	78.27 <sup>a</sup>	3.16 <sup>ab</sup>	0.04
Basal diet + Vitamin C + Vitamin E + Choline	132.22 <sup>b</sup>	52.72 <sup>c</sup>	78.12 <sup>a</sup>	3.21 <sup>ab</sup>	0.00
SEM	1.56	1.39	0.48	0.08	0.02
Significance	*	*	*	*	NS

Means having different superscripts in the same column are significantly ( $P \leq 0.05$ ) different

Means of feed conversion of the other experimental groups were comparable to that of the controls, with no significant differences among them. As given in Table 2, no significant differences were observed in mortality rates of turkey hens due to feeding the dietary treatment.

These results agree with the findings of Sahin *et al.* (2003) who reported that combined addition of dietary vitamin C and E, significantly increased feed intake of laying Japanese quails.

These results agree also with those of Wahyuni *et al.* (2011) who showed that vitamin E supplementation did not affect feed consumption in Kedu and Cemani hens. Krishnan and Scheideler (2010) showed that dietary supplementation with choline (choline chloride 60%) at 0, 500 and 1000 ppm did not show any difference in feed intake in 29-week-old White Leghorn hens. Panda *et al.* (2007) showed that daily feed consumption was not influenced by vitamin C supplementation at 200 mg/kg diet in White Leghorn layer. On the other hand, these results disagree with those of Shit *et al.* (2012) who showed that feed consumption of laying Japanese quails was significantly higher in groups fed L-ascorbic acid (250 and 500 ppm) supplemented diets than that of the control laying Japanese quails, at low ambient temperature. Ariana *et al.* (2011) found that feeding alpha-tocopheryl acetate diet (200 mg/kg) significantly improved feed intake in laying hens.

The present results agree also with those of Khan *et al.* (2011) who found that dietary supplementation of vitamin E improved egg production in laying hens. Ariana *et al.* (2011) found that feeding alpha -tocopheryl acetate diet (200 mg/kg) significantly improved egg production in laying hens. Noll (2000) indicated that egg production of turkey hens was improved by ascorbic acid supplementation (200 ppm) for 24 weeks. Sahin *et al.* (2003) showed that administration of vitamin C and E significantly increased egg production of laying Japanese quails.

The present results agree with those of Mbajjorgu (2011) who found that Venda hens supplemented with 200 and 500 mg of ascorbic acid per kg feed (Dry matter basis) produced heavier eggs than those produced by the unsupplemented birds. However, Heydari *et al.* (2009) found that dietary vitamin E (alpha -Tocopheryl acetate), 0 or 1000 mg/kg of diet had no significant effect on egg weight of laying hens, from 38-46 weeks of age.

The present results agree also with those of El-Zaiat *et al.* (2007) who showed that the addition of vitamin E to the diet (200 mg vitamin E/kg diet) caused an improvement of feed conversion of laying hens fed lead-polluted diet. Lin *et al.* (2004) found that the addition of 80 mg of vitamin E/kg diet gave the best feed conversion. Khan *et al.* (2011) found that dietary supplementation of vitamin E of heat-stressed poultry can improve feed efficiency.

In view of present results, laying hens fed diets containing vitamin C alone gave numerically better feed conversion ratio when compared with that of the control group. These results agree with those of Metwally (2004) who showed that supplemental vitamin C at the level of 2000 ppm improved feed conversion of Dandarawi laying hens.

The present results disagree with those of Çiftci *et al.* (2005) who found that mortality was significantly decreased in vitamin E-supplemented birds

(125 mg of alpha –tocopheryl acetate /kg of diet) compared to the controls. Chung *et al.* (2005) indicated that dietary supplementation with vitamin C to heat stressed Ross broiler breeder caused a significant decrease in mortality rate (15% - 20%).

**Egg Components and Egg Quality:**

Data on the effect of dietary single and combined addition of vitamin C, vitamin E and choline on egg components and certain parameters of egg quality of 42-week-old turkey hens are shown in (Table 3). The present results revealed that there were no significant differences among dietary treatments with respect to relative weights of egg albumen, egg yolk, egg yolk index, shell thickness or yolk color (Table 3). As indicated in Table 3, turkey hens fed diets supplemented with vitamin E alone or vitamin E plus choline produced eggs with lowered ( $P \leq 0.05$ ) percentages of egg shell than those of the control birds. Relative weights of egg shell of the other experimental groups were not significantly different from that of the control group.

**(Table 3): Effects of dietary single and combined addition of vitamin C , vitamin E and choline on egg quality and egg components of turkey hens at 42 weeks of age**

Items	Egg components (%)			Egg quality				
	Egg albumen %	Egg yolk %	Egg shell %	Egg shape index %	Egg yolk index %	Haugh units	Shell thickness (mm)	Yolk color
Control (Basal diet)	54.02	31.00	14.96 <sup>a</sup>	71.57 <sup>b</sup>	48.54	89.17 <sup>c</sup>	0.371	7.40
Basal diet +Vitamin C	56.30	30.06	13.62 <sup>ab</sup>	75.79 <sup>a</sup>	48.07	91.02 <sup>abc</sup>	0.397	7.30
Basal diet +Vitamin E	55.79	30.88	13.32 <sup>b</sup>	74.28 <sup>ab</sup>	49.42	95.58 <sup>ab</sup>	0.401	7.60
Basal diet + Choline	54.65	30.63	14.71 <sup>ab</sup>	75.62 <sup>a</sup>	48.37	91.11 <sup>abc</sup>	0.395	7.70
Basal diet +Vitamin C + Vitamin E	55.05	30.11	14.82 <sup>ab</sup>	75.38 <sup>a</sup>	48.36	89.62 <sup>c</sup>	0.388	7.40
Basal diet + Vitamin C + Choline	55.05	31.11	13.83 <sup>ab</sup>	75.44 <sup>a</sup>	48.45	90.40 <sup>c</sup>	0.388	7.40
Basal diet + Vitamin E + Choline	56.24	30.38	13.36 <sup>b</sup>	73.91 <sup>ab</sup>	47.82	96.49 <sup>a</sup>	0.388	7.80
Basal diet + Vitamin C + Vitamin E + Choline	55.20	30.82	13.96 <sup>ab</sup>	76.45 <sup>a</sup>	49.86	92.92 <sup>abc</sup>	0.398	7.60
SEM	0.93	0.82	0.48	1.13	1.30	1.74	0.957	0.27
Significance	NS	NS	*	*	NS	*	NS	NS

Means having different superscripts in the same column are significantly ( $P \leq 0.05$ ) different

The best means of egg shape index ( $p \leq 0.05$ ) were achieved by hens fed diets supplemented with a mixture of vitamin C, vitamin E and choline (76.45%), vitamin C alone (75.79%), choline (75.62%), vitamin C plus choline (75.44%) and vitamin C plus vitamin E (75.38%) in a descending order as compared to that of the control turkey hens. Egg shape indices of the other experimental groups were not significantly different from that of the controls. Compared with the control birds, turkey hens fed diets supplemented with vitamin E plus choline or with vitamin E alone produced eggs with better albumen quality as measured by Haugh units. No significant differences were observed in Haugh units among the other experimental groups of turkey hens.

The present results agree with those of Mohiti-Asli and Zaghari (2010) who found that shell thickness, yolk color and yolk weight of laying hens were not significantly affected by dietary supplementation with vitamin E. Radwan *et al.* (2008) showed that the addition of 200 mg vitamin E /kg of hen's diet numerically increased the percentage of egg shape index in El-Salaam strain laying hens. Kirunda *et al.* (2001) found that supplemental vitamin E can slightly improve egg albumen solids of laying hens.

The current results agree also with those of Heydari *et al.* (2009) who found that added dietary vitamin E (alpha -Tocopheryl acetate) at 1000 mg/kg had no significant effect on yolk weight, shell thickness of layers, from 38-46 weeks of age. Ebran and Bolukbasi (2011) found that dietary supplementation with 85 U vitamin E for laying hens increased Haugh units. Yan and Kim (2011) found that dietary supplemental vitamin E improved the Haugh units of laying hens compared with those of hens fed basal diet in 35-week-old (Hy-line brown) layers. Rama Rao *et al.* (2001) indicated that Haugh unit score and egg shell weight were not influenced by supplementary choline (760 and 1520 mg/kg) in diets of broiler breeders (29 to 48 weeks of age). Sahn and Sahn (2001) found that egg shell thickness and Haugh units of laying hens were improved due to feeding diet containing 250 mg vitamin C. Ajakaiye *et al.* (2011) found that Haugh Units of egg were higher in the vitamins C+E treated group (150 mg of L-ascorbic acid/kg of diet plus 150 mg of alpha-dl-tocopherol acetate/kg of diet) when compared to control group in laying hens at 39 weeks old.

On the other hand, the present results disagree with those of Ariana *et al.* (2011) who found that feeding alpha -tocopheryl acetate supplemented-diet (200 mg/kg) significantly improved yolk index in laying hens. Tsiagbe *et al.* (1988) observed an increase in yolk weight of eggs produced by hens given 1000 ppm of choline supplemented to a corn-soy based diet compared to that with no choline supplementation. Panda *et al.* (2008) reported that supplemental vitamin C to laying hens at 200 mg/kg diet significantly increased eggshell thickness. Çiftci *et al.* (2005) found a significant increase in relative weight of egg yolk in response to feeding vitamin E + vitamin C-supplemented diet (125 mg of alpha -tocopheryl acetate /kg of diet plus 200 mg of L-ascorbic acid of diet) as compared to the control group.



**Blood parameters of turkey hens :**

Data on the effect of dietary single and combined addition of vitamin C, vitamin E and choline on blood biochemical and haematological parameters of 48– week–old turkey hens are shown in (Table 4). As given in Table 4, concentrations of blood haemoglobin and serum total protein of the different experimental groups of turkey hens were comparable to that of the control birds, with no significant differences among them. Serum level of glucose for hens fed diets supplemented with vitamin C, choline, vitamin C plus vitamin E, vitamin C plus choline or a mixture of vitamin C, vitamin E and choline were significantly lower ( $P \leq 0.05$ ) than that of the controls. No significant differences were observed in serum glucose levels among the remaining experimental groups of turkey hens. Serum level of albumin for hens fed diets supplemented with choline alone or a mixture of vitamin C, vitamin E and choline were significantly lower ( $P \leq 0.05$ ) than that of the controls. No significant differences were observed in serum albumin levels among the remaining experimental groups of turkey hens.

**Table 4: Effects of dietary single and combined addition of vitamin C , vitamin E and choline on blood biochemical and hematological parameters on turkey hens at 48 weeks of age**

Items	Glucose (mg/dl)	Total protein (g/dl)	Albumin (g/dl)	Globulin (g/dl)	Total lipids (g/dl)	Cholesterol (mg/dl)	Hemoglobin (g/dl)	Haematocrit (%)
Control (Basal diet)	389.66 <sup>a</sup>	4.90	2.78 <sup>a</sup>	2.12 <sup>cd</sup>	3.57 <sup>a</sup>	303.00 <sup>a</sup>	10.33	36.63 <sup>ab</sup>
Basal diet + Vitamin C	347.33 <sup>bcd</sup>	5.83	2.73 <sup>ab</sup>	3.10 <sup>abc</sup>	2.41 <sup>b</sup>	274.00 <sup>b</sup>	11.50	31.00 <sup>d</sup>
Basal diet + Vitamin E	375.33 <sup>ab</sup>	4.53	2.60 <sup>ab</sup>	1.93 <sup>d</sup>	2.26 <sup>bc</sup>	198.33 <sup>d</sup>	10.73	35.80 <sup>abc</sup>
Basal diet + Choline	329.33 <sup>d</sup>	4.86	2.57 <sup>bc</sup>	2.29 <sup>bcd</sup>	2.46 <sup>b</sup>	282.33 <sup>ab</sup>	8.53	32.16 <sup>cd</sup>
Basal diet + Vitamin C + Vitamin E	341.33 <sup>cd</sup>	5.06	2.63 <sup>ab</sup>	2.43 <sup>bcd</sup>	1.79 <sup>c</sup>	169.33 <sup>e</sup>	11.23	35.06 <sup>abc</sup>
Basal diet + Vitamin C + Choline	353.00 <sup>bcd</sup>	5.90	2.74 <sup>ab</sup>	3.16 <sup>ab</sup>	3.32 <sup>a</sup>	227.33 <sup>c</sup>	10.16	34.00 <sup>bcd</sup>
Basal diet + Vitamin E+ Choline	368.33 <sup>abc</sup>	4.50	2.68 <sup>ab</sup>	1.82 <sup>d</sup>	2.49 <sup>b</sup>	220.66 <sup>cd</sup>	11.96	36.53 <sup>ab</sup>
Basal diet + Vitamin C + Vitamin E + Choline	353.00 <sup>bcd</sup>	5.90	2.40 <sup>c</sup>	3.50 <sup>a</sup>	3.36 <sup>a</sup>	258.33 <sup>b</sup>	12.10	38.20 <sup>a</sup>
SEM	9.47	0.33	0.05	0.31	0.18	8.48	0.60	1.22
Significance	*	NS	*	*	*	*	NS	*

Means having different superscripts at the same column are significantly ( $P \leq 0.05$ ) different

Serum globulin concentrations of hens fed the diets supplemented with vitamin C plus choline or a mixture of vitamin C, vitamin E and choline were significantly higher ( $P \leq 0.05$ ) than that of the control group, however, other experimental groups of hens exhibited insignificantly comparable levels of globulin to that of the controls. The turkey hens fed diets supplemented with vitamin C plus vitamin E, vitamin E, vitamin C, choline and vitamin E plus choline exhibited significantly lower ( $P \leq 0.05$ ) levels of serum total lipids compared with that of the control hens.

Serum levels of total lipids of hens fed diets supplemented with vitamin C plus choline or with the three feed additives were comparable to that of the control birds, with no significant differences among them. Dietary single and combined addition of vitamin C, vitamin E and choline caused significant reductions ( $P \leq 0.05$ ) in serum cholesterol levels as compared to that of the controls, with the exception of hens fed choline- supplemented diet which gave insignificantly comparable level of serum cholesterol. Dietary supplementation with vitamin C or choline resulted in significant decreases ( $P \leq 0.05$ ) in blood haematocrite values (%) of turkey hens compared with that of the controls, other groups exhibited comparable means of haematocrit value.

The present results agree with those of El-Medany (2008) who found that serum or plasma glucose content was significantly reduced when vitamin C (250 mg /kg diet) was added to the diet of laying hens. Sahin *et al.* (2003) showed that dietary supplementation with vitamin C and E either separately or in combination, at 250 mg/kg of both vitamin C and E decreased serum (plasma) glucose concentrations of laying Japanese quails. Imik *et al.* (2009) reported that the serum glucose level was lower in vitamin C group and vitamin E when compared with the control group given a normal diet without supplementation.

On the other hand, the present results disagree with the finding of Hassan *et al.* (2005) who found that choline supplementation (300 or 600 mg /kg feed) significantly increased blood serum level of albumin compared with that of the unsupplemented controls.

The present results agree with those of El-Medany (2008) who found that plasma levels of cholesterol and total lipids were significantly reduced when vitamin C (250 mg /kg diet ) was added to the diet of laying hens. Al-Daraji *et al.* (2001) indicated that levels of serum cholesterol were significantly decreased, as a result of ascorbic acid supplementation in both males and females of broiler breeders. Sahin *et al.* (2003) found that supplemental vitamin C and E either separately or in combination, (250 mg of L-ascorbic acid plus 250 mg alpha-tocopherol-acetate/kg of diet) decreased levels of serum (plasma) cholesterol of laying Japanese quails.

#### **Egg Fertility and Hatchability for turkeys:**

As presented in (Table 5), significantly higher ( $P \leq 0.05$ ) percentages of egg fertility and hatchability of total eggs were achieved by turkey birds fed diets supplemented with vitamin E alone, vitamin C plus choline, vitamin E plus choline or with a mixture of vitamin C, vitamin E and choline as compared to their control counterparts. The other experimental groups of birds exhibited insignificantly comparable means of egg fertility and total

hatchability compared to those of the control group. No significant differences were observed in hatchability of fertile eggs among the different experimental groups.

These results agree with those reported by Lin *et al.* (2004) who found that addition of 80 mg /kg vitamin E during the laying period increased egg fertility by 7.7% as compared with their controls. Hens fed 80 mg /Kg vitamin E exhibited an increase in percentage of hatchability by 13.4% as compared with those fed the unsupplemented diet . Sengul *et al.* (2008) showed that fertility rate and hatching rate improved with increasing levels of vitamin E at 30, 45, 60 and 75 mg/kg of laying partridges diets. Mbajorgu (2011) found that optimum ascorbic acid supplementation (750 mg kg DM feed) supported optimum hatchability in Venda laying hens, which was mainly attributable to its positive effect on egg weight and hatchability.

**Table 5: Effects of dietary single and combined addition of vitamin C , vitamin E and choline on egg fertility and hatchability of turkey hens during the experimental period**

Items	Egg Fertility (%)	Hatchability of total eggs (%)	Hatchability of fertile eggs (%)
Control (Basal diet)	93.72 <sup>d</sup>	86.13 <sup>c</sup>	91.87
Basal diet + Vitamin C	96.34 <sup>abcd</sup>	91.14 <sup>abc</sup>	94.56
Basal diet + Vitamin E	96.89 <sup>abc</sup>	92.06 <sup>ab</sup>	95.02
Basal diet + Choline	94.84 <sup>cd</sup>	86.97 <sup>c</sup>	91.53
Basal diet + Vitamin C +Vitamin E	95.86 <sup>cd</sup>	87.75 <sup>bc</sup>	91.70
Basal diet + Vitamin C + Choline	98.81 <sup>ab</sup>	91.77 <sup>ab</sup>	92.87
Basal diet + Vitamin E+ Choline	98.97 <sup>a</sup>	91.58 <sup>ab</sup>	92.53
Basal diet + Vitamin C + Vitamin E + Choline	98.50 <sup>ab</sup>	93.55 <sup>a</sup>	94.98
SEM	0.93	1.66	1.46
Significance	*	*	NS

Means having different superscripts at the same column are significantly ( $P \leq 0.05$ ) different

**Selected slaughter measurements:**

In turkey hens (Table 6), excepting for a significant increase in oviduct length of hens fed vitamin E-supplemented diet compared with the control group, the relative weights of their carcass, liver, heart, spleen, ovary and oviduct were not significantly affected by dietary treatments.

The present results agree with those of El-Zaiat *et al.* (2007) who found that the dietary addition of 200 mg vitamin E/kg diet) to Mandara laying hens insignificantly improved percentages of carcass, heart, spleen, and liver as compared to the control group . Rama Rao *et al.* (2001) reported that liver weight was not influenced by supplementary choline chloride (50%) and indicated that liver weight was not influenced by supplementary choline (760 and 1520 mg/kg) in broiler breeders (29 to 48 weeks of age). Fouladi *et al.* (2011) showed that dietary choline chloride supplement had no positive effect on meat yield and carcass weight in Japanese quails.

**Table 6: Effects of dietary single and combined addition of vitamin C , vitamin E and choline on some relative weights of carcass and some vital internal organs of 48-week-old turkey hens :**

Items	Carcass %	Liver %	Spleen %	Heart%	Ovary%	Oviduct%	Oviduct length(cm)
Control (Basal diet)	72.71	2.56	0.116	0.40	2.20	2.91	62.00 <sup>bc</sup>
Basal diet + Vitamin C	72.16	2.78	0.116	0.36	2.25	2.60	64.33 <sup>bc</sup>
Basal diet + Vitamin E	73.14	1.90	0.123	0.37 <sup>b</sup>	2.45	3.09	76.66 <sup>a</sup>
Basal diet + Choline	72.68	2.27	0.123	0.37	2.23	2.22	69.66 <sup>ab</sup>
Basal diet + Vitamin C +Vitamin E	73.26	2.35	0.110	0.38	2.00	1.93	55.00 <sup>c</sup>
Basal diet + Vitamin C + Choline	73.20	1.92	0.113	0.38	1.90	2.82	71.00 <sup>ab</sup>
Basal diet + Vitamin E + Choline	72.42	2.45	0.113	0.46	2.50	2.65	61.33 <sup>bc</sup>
Basal diet + Vitamin C + Vitamin E + Choline	71.90	3.16	0.113	0.38	2.16	2.59	65.66 <sup>b</sup>
SEM	0.92	0.28	0.003	0.02	0.32	0.40	3.05
Significance	NS	NS	NS	NS	NS	NS	*

Means having different superscripts at the same column are significantly (P≤0.05) different

In conclusion, dietary supplementation with vitamin E had positive effects on feed conversion, hen-day egg production rate (P≤0.05), egg fertility and total hatchability and length of oviduct of turkey hens as compared to those of the control hens.

## REFERENCES

- Ajakaiye, J. J. ; M . Cuesta-Mazorra and J.R Garcia-Diaz (2011). Vitamins C and E can alleviate adverse effects of heat stress on live weight and some egg quality profiles of layer hens. *Pak Vet J*, 31(1): 45-49.
- Al-Daraji, H. J.; E. H. Al-Mashhadani and A. K. Al-Athari (2001). Effect of ascorbic acid supplementation in the diets on haematological traits of Fawbro broiler breeders reared under hot climate. *Indian Journal of Animal Sciences*, 71 (9) : 857-859.
- Ariana, M.; A. Samie ; M. A. Edriss and R. Jahanian (2011). Effects of powder and extract form of green tea and marigold, and alpha tocopheryl acetate on performance, egg quality and egg yolk cholesterol levels of laying hens in late phase of production. *Journal of Medicinal Plants Research*, 5 (13) : 2710-2716.
- Azzi A. and A. Stocker (2000): Vitamin E: non-antioxidant roles. *Progress in Lipid Research*, 39 : 231–255.
- Chan, K. M. and E. A. Decker (1994). Endogenous skeletal muscle antioxidants. *Crit. Rev. Food Sci. Nutr*, 34: 403-426.
- Chan, M. M. (1984). Choline and carnitine. *Handbook of vitamins. Nutritional, Biochemical and Clinical Aspects*. L. J. Machlin ed. NY: Marcel Dekker, Inc. , chapter 14,pp.549 - 570.
- Chung, M. K. ; J. H. Choi ; Y. K. Chung and K. M. Chee (2005). Effects of dietary vitamins C and E on egg shell quality of broiler breeder hens exposed to heat stress. *Asian-Aust. J. Anim. Sci.*, 18 (4) : 545-551.
- Ciftci, M. ; O. Nihat Ertas and T. Guler (2005). Effects of vitamin E and vitamin C dietary supplementation on egg production and egg quality of laying hens exposed to a chronic heat stress. *Revue Méd. Vét.*, 156 (2) :107-111.
- Doumas, B.T. ; D. Watson and H. G. Biggs (1977). Albumin standards and the measurement of blood albumin with bromocrisol green. *Clin. Chem. Acta.*, 31: 87.
- Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics* ,11:1–42.
- Ehran, M. K. and S. C. Bolukbasi (2011). Effects of feeding diets supplemented with vitamin E and vitamin C on performance, egg quality and stereological and structural analysis of the liver of laying hens exposed to heat stress. *Italian Journal of Animal Science*, 10: (4) e58.
- Eisen, E. J. ; B. B. Bohren , and H. E. Mckeen (1962). The haugh unit as a measured of egg albumen quality. *Poult. Sci.*41: 1461-1468.
- El-Medany, N. M. (2008). Effect of dietary chromium and ascorbic acid supplementation on performance of laying hens under Egyptian summer conditions. *Egyptian Journal of Nutrition and Feeds*, 11: (1) 201-214.

- El-Zaiat, A. A. A.; S. A. M. Attia ; M. S. Shoeib and I. A. Azazi (2007). Laying hens performance and some histological changes as affected by the diet polluted with the lead added with clay or vitamin E. *Egyptian Poultry Science*, 27 (2) : 411-439.
- Fouladi, P. ; R . Salamat Doust Nobar ; A. Ahmadzade; H. Aghdam Shahriar and A. Agajanzade (2011). Effect of choline chloride supplement on the internal organs and carcass weight of Japanese quail . *Annals of Biological Research*, 2 (5): 485-491.
- Halliwell, B. and J. M. C. Gutteridge (1989) . *Free Radicals in Biology and Medicine*. 2nd ed. Oxford University Press, New York, NY .
- Hassan, R. A. ; Y. A. Attia and E.H. El-Ganzory (2005). Growth, carcass quality and serum constituents of slow growing chicks as effected by betaine addition to diets containing different levels of choline. *International Journal of Poultry Science*, 4 (11): 840-850.
- Henry, R. J. ; D.C. Cannon and J.W. Winkelman (1974). *Clinical Chemistry, Principles and Techniques*, 2nd ed. Harper and Ro.
- Heydari, A.; M. Eslami; H. Roshanfekar; M. Bojarpur and M. R. Ghorbani (2009). Effects of different levels of processed fat and vitamin E on laying hens performance and egg yolk cholesterol. *Research Journal of Biological Sciences*, 4 (11) :1162-1164.
- Hunsaker, W.G. (1969). Species and sex differences in the percentage of plasma trapped in packed cell volume determination in avian blood. *Poult. sci.*, 48 : 907-909.
- Imik, H.; S. Ozkanlar ; O. Kaynar and M. Koc (2009). Effects of vitamin E, C, and alpha -lipoic acid supplementation on the serum glucose, lipid profile and proteins in quails under heat stress. *Bulletin of the Veterinary Institute in Pulawy*, 53 (3) : 521-526.
- Kettunen, H. ; S. Peuranen ; K. Tiihonen and M.T. Saarinen (2001). Intestinal uptake of betaine *in vitro* and the distribution of methyl groups from betaine, choline and methionine in the body of broiler chicks. *Comp. Bioch. and Physiol. Part A*, 128: 269-278.
- Khan, R. U. ; S. Naz ; Z. Nikousefat ; V. Tufarelli ; M. Javdani ; N. Rana and V. Laudadio (2011). Effect of vitamin E in heat-stressed poultry. *World's Poultry Science Journal*, 67 (3) : 469-478.
- Kirunda, D.F.K. ; S. E. Scheideler and S.R McKee (2001). The efficacy of vitamin E (DL-alpha-tocopheryl acetate) supplementation in hen diets to alleviate egg quality deterioration associated with high temperature exposure. *Poultry Sci.*, 80: 1378-1383.
- Krishnan, P. and S. E. Scheideler (2010). Effects of dietary choline, folic acid and vitamin B12 on laying hen performance, egg components and egg phospholipid composition" (2010). D. S.C. thesis. Univ. of Nebraska – Lincoln. Lincoln, NE.
- Lin, Y.F. ; S.J. Chang and A.L. Hsu (2004). Effects of supplemental vitamin E during the laying period on the reproductive performance of Taiwan native chickens. *Br. Poult. Sci.*, 45: 807-814.

- Mbajjorgu, C. A. (2011). Effect of dietary ascorbic acid supplementation on egg production, egg quality and hatchability of indigenous Venda chicken hens. *Asian Journal of Animal and Veterinary Advances*, 6 (8) : 823-830.
- McDowell, L. R. (1989). *Vitamins in Animal Nutrition*. Publ. Academic Press, N.Y. Mitchell, A. D., and N. J. Benevenga. 1978. The role of transamination in methionine oxidation in the rat. *J. Nutr.* , 108: 67-78.
- Metwally, M. A. (2004). Effect of dietary vitamin C on the performance of Dandarawi laying hens subjected to cold stress. *Egyptian Poultry Science*, 24 (2) : 465-481.
- Mohiti-Asli; M. and M. Zaghari (2010). Does dietary vitamin E or C decrease egg yolk cholesterol? *Biological Trace Element Research*. 138 (1/3) : 60-68.
- National Research Council (NRC) (1994). *Nutrient Requirements of Poultry*. 9th revised edition. National Academy Press. Washington, D.C.
- Noll, S. L.(2000). Vitamin C supplementation in turkey breeder diets. *Zootecnica International*,23: (11) 44-47.
- Panda, A. K. ; S. V. Ramarao ; M. V. L. N. Raju and R. N. Chatterjee (2008). Effect of dietary supplementation with vitamins E and C on production performance, immune responses and antioxidant status of White Leghorn layers under tropical summer conditions. *British Poultry Science*. 49: (5) 592-599.
- Panda, A. K. ; S. V. Ramarao and M. V. L. N. Raju (2007). Effect of vitamin C supplementation on performance, immune response and antioxidant status of heat stressed White Leghorn layers. *Indian Journal of Poultry Science*, 42 (2) :169-173.
- Pilaski, J. (1972). Vergleichende untersuchungen uberden hemoglobin halt des huhnner and geschlecht. *Arch Gelfuglek*, 36: 70-77.
- Puthongsiripon, U. (1998). Effect of strain and dietary vitamin E on hen performance, immune and antioxidant status during heat stress. M. S.C. thesis. Univ. of Nebraska – Lincoln. Lincoln, NE.
- Radwan Nadia, L. ; R.A. Hassan ; E.M. Qota and H.M. Fayek (2008). Effect of Natural Antioxidant on Oxidative Stability of Eggs and Productive and Reproductive Performance of Laying Hens. *International Journal of Poultry Science* 7 (2): 134-150.
- Rama-Rao, S. V. ; G. S. Sunder ; M. R. Reddy ; N. K. Praharaj ; M. V. Raju and A. K. Panda (2001).Effect of supplementary choline on the performance of broiler breeders fed on different energy sources. *Br Poult Sci. Jul*;42 (3): 362-367.
- Sahin, K.; N. Sahin ; M. Onderci ; M. F. Gursu and M. Issi (2003). Vitamin C and E can alleviate negative effects of heat stress in Japanese quails. *Journal of Food, Agriculture & Environment*. 1 (2) :244-249.
- Sahn, N. and K. Sahn (2001). Optimal dietary concentrations of vitamin C and chromium picolinate for alleviating the effect of low ambient temperature (6.2 degrees C) on egg production, some egg characteristics, and nutrient digestibility in laying hens. *Veterinari Medicina*, 46: (9/10) 229-236.

- SAS (2003). SAS User's Guide: Statistics. Version 8.2, SAS Institute Inc., Cary, NC.
- Sengul, T. ; M. Cetn ; B. Sogut and S. Yurtseven (2008). Effects of dietary vitamin E levels on egg production traits, fertility and hatching rate in laying partridges. *Journal of Applied Animal Research.* , 34 (1) :77-80.
- Sheard, N. F. and S. H. Zeisel (1989). Choline: an essential dietary nutrient? *Nutrition*, 5: 1-5
- Shit, N. A ; R. P. Singh ; A. K. V. H. Sastry ; R. Agarwal ; R Singh ; N. K. Pandey and J. Mohan (2012). Effect of dietary l-ascorbic acid (L-AA) on production performance, egg quality traits and fertility in Japanese quail (*Coturnix japonica*) at low ambient temperature. *Asian-Aust. J. Anim. Sci.* (25) , 7 : 1009 - 1014 .
- Tanaka K. ; T. Hashimoto ; S. Tokumaru ; H. Iguchi and S. Kojo (1997). Interactions between vitamin C and vitamin E are observed in tissues of inherently scorbutic rats. *Journal Nutrition*, 127: 2060–2064.
- Trinder, P. (1969). Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Ann. Clin. Biochem*,6: 24-27.
- Tsiagbe, V. K. ; M. E. Cook ; A. E. Harper, and M. L. Sunde (1988). Alterations in phospholipid composition of egg yolks from laying hens fed choline and methionine supplemented diets. *Poultry Sci.*, 67: 1717-1724.
- Wahyuni, H. I. ; N. Suthama ; I. Mangisah and T. A. Sarjana (2011). Egg quality and hatchability of reared Kedu and Cemani hens fed diet of farmer formulation supplemented with vitamin E. *Journal of the Indonesian Tropical Animal Agriculture.* 36 (1) : 61-68.
- Wang, Y. H. ; J. Leibholz ; W. L. Bryden and D. R. Fraser (1996). Lipid-peroxidation status as an index to evaluate the influence of dietary fats on vitamin E requirements of young pigs. *Br. J. Nutr.* , 75: 81-95.
- Watson, D. (1960). A simple method for the determination of serum cholesterol. *Clin. Chem. Acta*, 5: 637.
- Yan, L. and I. H. Kim (2011). Evaluation of natural and synthetic vitamin E on production performance, egg quality and vitamin E content of yolk in laying hens. *Journal of Applied Animal Research.* 39 (4) :381-385.
- Zollner, N. and K. Kirsch (1962). Test combination of the total lipids. *Zges.Exp.Med.*, 535-545 .



تأثير بعض مضادات الاكسدة والمواد المعطية لمجموعة الميثيل على الأداء الانتاجي والتناسلي لإناث الرومي  
عبد البصير حمزة محمد ريا<sup>1</sup> ، فوزى صديق عبدالفتاح اسماعيل<sup>1</sup> ، محيي الدين يوسف مصطفى<sup>2</sup> و مايكل عادل لبيب جورجي<sup>2</sup>  
1- قسم انتاج الدواجن - كلية الزراعة - جامعة المنصورة .  
2- معهد بحوث الانتاج الحيواني - مركز البحوث الزراعية - الدقى - الجيزة - مصر.

الهدف من هذه الدراسة هو تقييم تأثير استخدام بعض مضادات الاكسدة والمواد المعطية لمجموعة الميثيل على الاداء الانتاجي والتناسلي للرومي . أجريت هذه الدراسة فى محطة بحوث الانتاج الحيواني بمحلله موسى ، كفر الشيخ التابعة لمعهد بحوث الانتاج الحيواني ، مركز البحوث الزراعية ، وزارة الزراعة ، مصر .

استخدم فى هذه الدراسه عدد 168 طائر (انثى) من سلالة الرومي البرونز ، عمر 36 اسبوع . وزعت الطيور بطريقتة عشوائية الى 8 معاملات تجريبية ، احتوت كل معاملة على 3 مكررات متساوية . تم تسكين الطيور بطريقتة فردية فى بطاريات . استغرقت التجربة 12 اسبوع من عمر 36-48 اسبوع .

غذيت الطيور بعليقة اساسية (كنترول) مكونة من ذرة صفراء وكسب فول صويا مضاف اليها فيتامين C بمعدل 200 مجم/كجم علف ، فيتامين E بمعدل 250 مجم/كجم علف (مضادات اكسدة) والكولين بمعدل 2.25 جم/كجم علف (مصدر لمجموعة الميثيل) بصورة مفردة أو مجمعة . وكان أهم النتائج المتحصل عليها كالتالى :-

- أعطت المعاملة المضاف اليها فيتامين E تأثيرات ايجابية على معدل التحويل الغذائى ، معدل انتاج البيض اليومى ( $P \leq 0.05$ ) ، نسبة خصوبة البيض ، نسبة الفقس الكلية وطول قناة البيض لاناث الرومي مقارنة بالاناث فى مجموعة الكنترول .

- لم تكن هناك فروق معنوية بين مختلف المعاملات التجريبية فى معدل النفوق ، والوزن النسبى لكل من البياض والصفار ، دليل الصفار ، سمك القشرة ، لون الصفار ، تركيزات كل من الهيموجلوبين والبروتينات الكلية فى بلازما الدم ، نسبة الفقس للبيض المخصب ومواصفات الذبيحة لاناث الرومي .

يستنتج من النتائج المتحصل عليها ان اضافة فيتامين E الى العليقة بمعدل 250 مجم/كجم يمكن ان يحدث تأثيرات ايجابية على الاداء الانتاجي والتناسلي لإناث الرومي .

قام بتحكيم البحث

أ.د / خليل الشحات شريف

أ.د / عبد الله على غزاله

كلية الزراعة - جامعة المنصورة

كلية الزراعة - جامعة القاهرة