

## **MODULATION OF AFLATOXINS TOXICITY IN RATS BY ETHANOLIC POMEGRANATE PEEL (*Punica granatum.L*) EXTRACT**

Tohame, Amany A.\*; Amel A. Abo-Hagger\*\* and Y. M. Abd El-Shafea\*\*

\* Department of Zoology and Entomology, Faculty of Science, Helwan University, Helwan, Egypt

\*\* Regional Center for Foods and Feeds (RCFF), Agricultural Research Center (ARC), Giza, Egypt

### **ABSTRACT**

This work was conducted to investigate the effect of ethanolic pomegranate peels (PP) extract on liver and kidney structure and function in rats experimentally exposed to different doses of aflatoxins (AFS). An experiment was conducted for 6 weeks, 36 male rats were divided into 6 groups (6 rats in each). G<sub>1</sub> served as vehicle treated control, G<sub>2</sub> received PP extract via gavage (50 mg/kg body weight (b.w.) daily). G<sub>3</sub> received AFS via gavage (1 mg/kg b.w.) twice/week. G<sub>4</sub> received AFS via gavage (2 mg/kg b.w.) twice/week. G<sub>5</sub> received AFS twice/week (1 mg/kg b.w.) + (PP) extract daily (50 mg/kg b.w.). G<sub>6</sub> received AFS twice/week (2 mg/kg b.w.) + (PP) extract daily (50 mg/kg b.w.). Liver enzymes (Alanine Amino Transferase (ALT), Aspartate Amino Transferase (AST), Alkaline Phosphatase (ALP)), Total Protein (TP), albumin and renal biochemical parameters (creatinine, urea and uric acid) were estimated in serum at 3<sup>rd</sup> and 6<sup>th</sup> week post treatment, while pathological examination of liver and kidney was performed at the end of experiment. Result demonstrated that both aflatoxins administrated levels caused elevation in AST, ALT, ALP, creatinine, urea and uric acid and a decrease in TP and albumin in dose and time dependant manner, while synergistic administration of PP extract and AFS resulted in limitation of the negative effects of AFS on the tested parameters especially with low AFS dose and longer time. Pathological examination revealed that the severity of damage in liver and kidney correlated with aflatoxins' doses. While PP extract administration was able to protect liver and kidney against low AFS dose and minimize the damage of liver and kidney induced by high AFS dose.

**Keywords:** Aflatoxin, Pomegranate, Pathological examination, Liver enzymes and renal biochemical parameters

### **INTRODUCTION**

Aflatoxins (B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub>) are a group of toxic compounds produced by toxigenic strains of *Aspergillus flavus* and *Aspergillus parasiticus*. These fungi can grow on a wide variety of food commodities (Wilson and Payne, 1993). Humans are exposed to AFS directly by the consumption of contaminated food or indirectly by the consumption of animal products (i.e., fish, meat, milk and eggs) derived from animals that consumed AFS contaminated feed (Abdelhamid *et al.*, 1994 and Bennett and Klich, 2003). AFS are well known as hepatotoxic, hepatocarcinogenic and mutagenic agents. These effects are mainly due to adduct formation with DNA, RNA and protein. In addition, it also causes lipid peroxidation as well as oxidative damage to DNA. AFB<sub>1</sub> possess genotoxic potential in a variety of test systems. Other aflatoxin has not been so extensively investigated, but in

a variety of studies B<sub>2</sub>, G<sub>1</sub>, G<sub>2</sub> and M<sub>1</sub> have all shown evidence of genotoxicity (Verma, 2004). The AFS present in food and feed are hazardous to both human and animal health (Do and Choi, 2007). Therefore, development and application of intervention strategies that are highly effective and economically feasible against dietary AFB<sub>1</sub> contamination and aflatoxicosis are critical for improving human and animal health, especially in high-risk areas.

AFS toxicity may ensue through the generation of intracellular reactive oxygen species (ROS) like superoxide anion, hydroxyl radical and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) during the metabolic processing of AFB<sub>1</sub> by cytochrome P450 in the liver (Preetha *et al.*, 2006).

Ueno *et al.* (1991) indicated that active oxygen scavengers reduce mutation induced by various mutagens. It has been suggested that compounds which possess antioxidants activity can inhibit mutation and cancer because they can scavenge a free radicals or induce antioxidants enzymes (Hochstein and Atallah, 1988). Dietary intake of natural antioxidants could be an important factor in body's defense mechanism against many mutagens and carcinogens, also many antioxidants are being identified as anticarcinogens. Many plant polyphenols, have been shown to act as potent antimutagenic and anticarcinogenic agents (Tanaka *et al.*, 1993 and Yen and Chen, 1994).

The pomegranate, *Punica granatum* L., is an ancient, mystical, unique fruit borne on a small, long-living tree cultivated throughout the Mediterranean region, as far north as the Himalayas, in Southeast Asia, and in California and Arizona in the United States. In addition to its ancient historical uses, pomegranate is used in several systems of medicine for a variety of ailments. The synergistic action of the pomegranate constituents appears to be superior to that of single constituents. In the past decade, numerous studies on the antioxidant, anticarcinogenic, and anti-inflammatory properties of pomegranate constituents have been published, focusing on treatment and prevention of cancer, cardiovascular disease, diabetes, dental conditions, erectile dysfunction, bacterial infections and antibiotic resistance, and ultraviolet radiation-induced skin damage. Other potential applications include infant brain ischemia, male infertility, Alzheimer's disease, arthritis, and obesity (Julie, 2008). Negi *et al.* (2003) showed that the pomegranate peel (PP) extracts have both antioxidant and antimutagenic properties and may be exploited as biopreservatives in food applications and nutraceuticals. PP extract contains substantial amounts of polyphenols such as ellagic tannins, ellagic acid and gallic acid. It has been used in preparation of tinctures, cosmetics, therapeutic formulae and food recipes (Nasr *et al.*, 1996). Guo *et al.* (2003) found that PP had highest antioxidant activity among the peel, pulp and seed fractions of 28 kinds of fruits commonly consumed in China. Moreover, Li *et al.* (2006) concluded that PP extract appears to have more potential as health supplement rich in natural antioxidants than the pomegranate pulp extract.

In this study, we investigated the modulation of ethanolic PP extract on toxic effects of AFS in male Albino rats.

## MATERIALS AND METHODS

### Chemicals

All chemicals were of analytical grade. Chemicals were purchased from Sigma Chemical Co. (St Louis, MO, USA) and Solvents were purchased from Merck (Darmstadt, Germany).

### Microorganism

*Aspergillus flavus* NRRL (3145) was obtained from the National Research Center (Dokki, Giza).

### Standard Aflatoxins

Aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> were obtained from Sigma Chemical Company (St. Louis, Mo USA). The purity of these materials were checked by thin layer chromatography [silica gel aluminium sheets 20x20 cm purchased from Merck (Darmstadt, Germany) using a mixture of diethyl ether: methanol: distilled water (96: 3: 1, v/v/v) as a running solution] and each gave one spot.

### Aflatoxin production and assays

#### Preparation of media for *Aspergillus flavus* growth

**a) Potato-Dextrose-Agar Media (PDA):** This medium was prepared according to *Shotwell et al. (1966)* as follows:

Flask 1 contained: Dextrose (20g), CaCO<sub>3</sub> (0.2g), MgSO<sub>4</sub>.7 H<sub>2</sub>O (0.2g) and distilled water (100 ml)

Flask 2 contained: Agar (15g) and distilled water (400ml)

Flask 3 contained: Potato (Spilled and sliced) (200g) and distilled water (500ml)

The contents of flask 3 were brought shortly to 121 °C in autoclave and filtered through cheesecloth. The solution was brought up to original volume simultaneously. The agar in flask 2 was melted and the solution in flask 1 was heated to boiling. Contents of the three flasks were mixed together and autoclaved for 15 min. at 121 °C.

**b) Yeast Extract Sucrose Media (YES):** This medium yields high amount of aflatoxin especially B<sub>1</sub> and G<sub>1</sub> with *A. flavus*. It was prepared according to method of *Davis et al. (1966)* as follows: Yeast extracts (20g), Sucrose (200g), FeSO<sub>4</sub>.7H<sub>2</sub>O (10mg), ZnSO<sub>4</sub>.7H<sub>2</sub>O (5mg), MnSO<sub>4</sub>.4H<sub>2</sub>O (1mg) and distilled water (1000ml) were dissolved, mixed and autoclaved for 15 min at 121°C.

### Fungal growth

Inoculum was prepared by incubating PDA slant tubes (1.5x15 cm) inoculated with *A. flavus* spores for 7 to 21 days at 28 °C. Spores of 10-day old (*A. flavus* culture) were scrapped loose with a loop after adding 3 ml sterile distilled water to each slant (*Shotwell et al., 1966*). The Spores scraped by adding sterile distilled water to the surface growth on agar slant and an aliquot amount from the resulting spore suspension (1 ml) was added to conical flasks (2 liters) containing yeast extract media. Mycelial mats after 10-day incubation at 30°C were broken with a glass rod and collected by filtration through filter paper. Culture filtrates were extracted with chloroform (1:2, v/v). (*Abd El-Mageed, 1987*). The chloroform extract was evaporated

until obtaining dry film in rotary evaporator. Then the obtained dry film containing AFS was reconstituted with dimethyl sulfoxide (DMSO) and used for quantitative analysis of AFS and then preparation of AFS doses.

#### **Determination of aflatoxins concentration in solution**

The concentration of AFS in DMSO solution was determined using HPLC technique (Agilent 1100 Series U.S.A with column C<sub>18</sub>, Lichrospher 100 RP-18, 5µm×25cm) as follows: The mobile phase consisted of water: methanol: acetonitrile (54:29:17, v/v/v) at a flow rate of 1ml/min. The excitation and emission wavelengths for all aflatoxins were 362 and 460nm (Fluorescence detector), respectively (Roos *et al.*, 1997).

#### **Preparation of aflatoxins dose**

The DMSO containing AFS solution consisted of a mixture of aflatoxin B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> at a total concentration of 34mg AFS/ ml as a ratio of 8: 2: 4: 1, respectively used to prepare 2 final concentrations of AFS, 1mg AFS/5ml DMSO and 2mg AFS/5ml DMSO.

#### **Pomegranate peels preparation and extraction.**

Peel was obtained from juice stores. It was air dried and milled using laboratory mill to pass a 1.0 mm-size, and then dried again in a cabinet oven with air circulation at 60°C for 16h. Then kept in refrigerator prior to extraction.

PP was extracted according to the method of Panovska *et al.* (2005). Ten g of dry powder were extracted with 100 ml of 70% ethanol in a screw-capped flask and shaken at room temperature for 24 h. The extracts were centrifuged at 5000 rpm for 10 min while the residue was re-extracted under the same conditions twice and filtered through a Büchner funnel with filter paper (Whatman No.1). The 70% ethanol extracts were concentrated under reduced pressure, lyophilized to obtain powders, and stored at 4 °C.

#### **Animal and treatment**

Thirty-six male Albino rats weighing 110±5g (provided by the Laboratory Animal Center, Faculty of Veterinary Medicine, Cairo University) were housed in stainless steel cages in animal house in Regional Center for Foods and Feeds, Agricultural Research Center, Ministry of Agriculture, Giza, Egypt under controlled light and temperature conditions 12/12 hours light: dark cycle, 22-33 °C). During the acclimation period (1 week) and experimental period (6 weeks), a basal diet and tap water were supplied *ad libitum*. The basal diet consisted of casein 20%, corn oil 10%, wooden fibers 5%, salt mixture 4%, vitamin mixture 1% and starch 60% (Lane – Peter and Pearson, 1971). Rats were divided into six groups (G) and treated for 6 weeks as follows: G<sub>1</sub>: served as vehicles treated control (received 5ml DMSO/ kg b.w. daily + 5ml DMSO/ kg b.w. twice/ week on Monday and Thursday, orally by gavage); G<sub>2</sub>: received orally 50mg PP extract powder/ kg b.w. suspended in 5ml DMSO daily + 5ml DMSO/ kg b.w. twice/week on Monday and Thursday; G<sub>3</sub>: received orally low dose of AFS (1mg AFS suspended in 5ml DMSO/ kg b.w.) twice/ week on Monday and Thursday + 5ml DMSO/ kg b.w. daily; G<sub>4</sub>: received orally high dose of AFS (2mg AFS suspended in 5ml DMSO/ kg b.w.) twice/ week on Monday and Thursday + 5ml DMSO/ kg b.w. daily; G<sub>5</sub>: received orally low dose of AFS (1mg AFS suspended in 5ml DMSO/ kg b.w.) twice/ week on Monday and Thursday + 50mg PP extract powder

suspended in 5ml DMSO/ kg b.w. daily; and G<sub>6</sub>: received orally high dose of AFS (2mg AFS suspended in 5ml DMSO/ kg b.w.) twice/ week on Monday and Thursday + 50mg PP extract powder suspended in 5ml DMSO/ kg b.w. daily.

At the 3<sup>rd</sup> week and 6<sup>th</sup> week of the treatment period, blood samples were collected from the retro-orbital venous plexus under light CO<sub>2</sub> anesthesia and the sera were separated and stored at -20 °C until analysis. Rats were scarified at the end of experiment and liver and kidney were removed for histological examination.

#### **Biochemical assays**

The Alera Clinical Chemistry (ACE) System Automatic Analyzer (Alfa Wasserman Corporation) was employed to measure the following parameters: aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, total protein, albumin, creatinine, urea and uric acid.

#### **Histopathological techniques**

Samples for histopathological examination were fixed in 10% neutralized formalin for 2 days, washed in tap water, dehydrated in ascending grades of ethyl alcohol and finally cleared with xylene and embedded in paraffin wax. The paraffin blocks were five – micron cutted and stained by Haematoxylin and Eosin as described by Pearse (1968).

#### **Statistical analysis**

The data obtained in the present work represented in tables as mean ± standard error. Statistical analysis for the collected data was done according to the procedure outlined by Gomez and Gomez (1984). The treatment means were compared using the least significant difference test (LSD) at the 5% level of probability as outlined by Waller and Duncan (1969) using the Duncan test institute program.

## **RESULTS**

### **Biochemical parameters**

#### **Liver function**

The *in vivo* study revealed that rats administrated AFS orally (G<sub>3</sub> and G<sub>4</sub>) significantly increased the serum activity of ALT, AST and ALP but decreased the TP and albumin levels in dose and time dependant manner when compared with control ( $P<0.05$ ) (Table 1). While no significant differences were found in their levels in serum of rats administrated with PP extract (G<sub>2</sub>) comparing with control one ( $P<0.05$ ) in both experimental times. Simultaneous administration of rats with PP extract and different doses of AFS (G<sub>5</sub> and G<sub>6</sub>) resulted in reduction of elevated levels of ALT, AST and ALP, elevation of depleted level of TP and restore level of albumin to normal (comparing to groups treated by AFS alone and control group). The ameliorating effect of PP extract on liver function was time dependent and more obvious on low AFS dose.

**Table (1): Activity of ALT, AST and ALP and concentration of TP and albumin in serum of rats during experimental periods (means ± SE).**

Parameters	ALT (U/L)		AST (U/L)		ALP (U/L)		TP (mg/dl)		Albumin (mg/dl)	
	3 weeks	6 weeks	3 weeks	6 weeks	3 weeks	6 weeks	3 weeks	6 weeks	3 weeks	6 weeks
G <sub>1</sub>	52.3± 0.89 <sup>g</sup>	60.3± 3.38 <sup>g</sup>	87.0± 4.36 <sup>h</sup>	107.3± 2.19 <sup>gh</sup>	349.3± 19.67 <sup>h</sup>	524.30± 15.07 <sup>g</sup>	7.84± 0.05 <sup>ab</sup>	8.09± 0.09 <sup>a</sup>	3.85± 0.07 <sup>ab</sup>	3.98± 0.12 <sup>a</sup>
G <sub>2</sub>	54.0± 2.08 <sup>g</sup>	59.0± 2.64 <sup>fg</sup>	90.67± 3.18 <sup>gh</sup>	112.3± 5.04 <sup>fg</sup>	368.0± 9.45 <sup>h</sup>	534.3± 17.14 <sup>g</sup>	7.93± 0.17 <sup>ab</sup>	8.15± 0.25 <sup>a</sup>	3.85± 0.06 <sup>ab</sup>	3.90± 0.10 <sup>ab</sup>
G <sub>3</sub>	126.3± 10.36 <sup>c</sup>	140.67± 10.27 <sup>c</sup>	189.67± 5.78 <sup>c</sup>	236.3± 16.3 <sup>b</sup>	668.0± 11.01 <sup>d</sup>	847.30± 24.34 <sup>b</sup>	6.98± 0.24 <sup>de</sup>	6.74± 0.29 <sup>e</sup>	3.63± 0.13 <sup>b</sup>	3.62± 0.06 <sup>b</sup>
G <sub>4</sub>	164.67± 12.34 <sup>b</sup>	189.67± 5.78 <sup>a</sup>	246.0± 8.62 <sup>b</sup>	306.0± 12.2 <sup>a</sup>	785.3± 21.18 <sup>c</sup>	1020± 30.61 <sup>a</sup>	6.84± 0.06 <sup>e</sup>	6.62± 0.25 <sup>e</sup>	3.30± 0.17 <sup>c</sup>	3.27± 0.10 <sup>c</sup>
G <sub>5</sub>	84.0± 3.79 <sup>de</sup>	77.0± 1.52 <sup>ef</sup>	160.0± 2.65 <sup>d</sup>	121.67± 4.63 <sup>ef</sup>	587.0± 5.86 <sup>ef</sup>	544.3± 23.88 <sup>g</sup>	7.47± 0.17 <sup>bcd</sup>	7.67± 0.17 <sup>abc</sup>	3.75± 0.04 <sup>ab</sup>	3.80± 0.05 <sup>ab</sup>
G <sub>6</sub>	98.0± 5.55 <sup>d</sup>	86.3± 1.76 <sup>de</sup>	198.67± 6.44 <sup>c</sup>	142.3± 6.23 <sup>de</sup>	631.3± 3.71 <sup>de</sup>	591.0± 15.17 <sup>ef</sup>	7.20± 0.20 <sup>cde</sup>	7.55± 0.14 <sup>a-d</sup>	3.73± 0.16 <sup>ab</sup>	3.68± 0.06 <sup>ab</sup>
LSD <sub>0.05</sub>	18.353		22.17		51.45		0.55		0.3	

The various superscript letters in each parameter indicate statistically significant differences in the Duncan test, at *P* <0.05

### Kidney function

The current results presented in Table 2 clearly indicated that rats treated by different doses of AFS showed significant increase in serum levels of creatinine, urea and uric acid comparing to that of control group (*P*<0.05). The increase of their levels was parallel to dose of AFS and time of administration. While there were no changes in their levels in serum of group treated by PP extract comparing to control group (*P*<0.05). The co-administration of PP extract and AFS in G<sub>5</sub> and G<sub>6</sub> rats resulted in reduction in increased levels of creatinine, urea and uric acid in serum in both times of treatment (comparing to groups treated by AFS alone and control group).

**Table (2): Creatinine, urea and uric acid concentration in serum of rats during experimental periods (means ± SE).**

Parameters	Creatinine(mg/dl)		Urea(mg/dl)		Uric acid(mg/dl)	
	3weeks	6weeks	3weeks	6weeks	3weeks	6weeks
G <sub>1</sub>	0.42± 0.01 <sup>g</sup>	0.47± 0.01 <sup>g</sup>	40± 1.52 <sup>g</sup>	51.3± 1.20 <sup>ef</sup>	1.26± 0.04 <sup>f</sup>	1.36± 0.06 <sup>ef</sup>
G <sub>2</sub>	0.45± 0.01 <sup>g</sup>	0.49± 0.03 <sup>fg</sup>	42.3± 1.76 <sup>g</sup>	52.3± 0.33 <sup>ef</sup>	1.28± 0.04 <sup>f</sup>	1.32± 0.07 <sup>ef</sup>
G <sub>3</sub>	0.81± 0.02 <sup>d</sup>	0.99± 0.04 <sup>c</sup>	66.0± 2.30 <sup>c</sup>	80.0± 2.08 <sup>b</sup>	2.19± 0.12 <sup>c</sup>	2.80± 0.21 <sup>b</sup>
G <sub>4</sub>	1.14± 0.10 <sup>b</sup>	1.33± 0.11 <sup>a</sup>	84.0± 2.08 <sup>b</sup>	99.0± 2.51 <sup>a</sup>	3.02± 0.08 <sup>b</sup>	4.16± 0.03 <sup>a</sup>
G <sub>5</sub>	0.63± 0.01 <sup>ef</sup>	0.54± 0.01 <sup>efg</sup>	56.0± 2.08 <sup>de</sup>	49.0± 2.08 <sup>f</sup>	1.71± 0.13 <sup>d</sup>	1.63± 0.09 <sup>de</sup>
G <sub>6</sub>	0.82± 0.03 <sup>d</sup>	0.65± 0.02 <sup>e</sup>	65.3± 2.85 <sup>c</sup>	59.3± 1.45 <sup>c</sup>	1.85± 0.03 <sup>d</sup>	1.74± 0.13 <sup>d</sup>
LSD <sub>0.05</sub>	0.143		5.73		0.299	

The various superscript letters in each parameter indicate statistically significant differences in the Duncan test, at *P* <0.05

The major reduction in their levels was noticed at 6 week of administration and with low AFS dose. No significant differences were noticed in ability of PP extract to reduce toxic effect of high dose of AFS on urea and uric acid between 3<sup>rd</sup> and 6<sup>th</sup> week of experiment.


### **Histopathological examination**

#### **Liver**


Microscopical examination of rat's liver from G<sub>1</sub> (vehicle treated control) revealed the normal marked histological structure of hepatic lobule (Fig.1). Also no histological changes were noticed in liver of rat from G<sub>2</sub> which treated with PP extract (Fig. 2).

Conversely, the examined sections from G<sub>3</sub> which treated by low AFS dose showed vacuolar degeneration of hepatocytes (Fig. 3), kupffer cells activation, individual cell necrosis of sporadic hepatocytes (Fig. 4) and focal hepatic necrosis (Fig. 5). Moreover, liver of rat of G<sub>4</sub> which treated with high AFS dose showed dramatic changes. Liver showed karyomegaly of some nuclei, apoptosis of hepatocytes (Fig. 6), portal infiltration with leucocytes (Fig. 7), cytokaryomegaly of hepatocytes, oval cells proliferation (Fig. 8) and mitotic figures (Fig. 9).


Non significant marked histopathological changes were noticed in liver of rat of G<sub>5</sub> (Fig. 10) which treated with low AFS dose and PP extract. Meanwhile, liver of G<sub>6</sub> (treated by high AFS dose and PP extract) showed kupffer cell proliferation and necrosis of sporadic hepatocytes (Fig. 11).




**Fig. (1):** Section in liver of rat from G<sub>1</sub> showing the normal structure of hepatic lobule (H and E X 200).




**Fig. (2):** Section in liver of rat from G<sub>2</sub> showing non significant histopathological changes (H and E X 200).




**Fig. (3):** Section in liver of rat from G<sub>3</sub> showing vacuolations of hepatocytes (H and E X 200).




**Fig. (4):** Section in liver of rat from G<sub>3</sub> showing kupffer cells activation and necrosis of sporadic hepatocytes (H and E X 200).




**Fig. (5):** Section in liver of rat from G<sub>3</sub> showing focal hepatic necrosis (H and E X 200).




**Fig. (6):** Section in liver of rat from G<sub>4</sub> showing karyomegaly of some nuclei and apoptosis of hepatocytes (H and E X 200).



**Fig. (7):** Section in liver of rat from G<sub>4</sub> showing portal infiltration with leucocytes (H and E X 200).




**Fig. (8):** Section in liver of rat from G<sub>4</sub> showing cytokaryomegaly of hepatocytes associated with oval cell proliferatin (H and E X 200).



**Fig. (9):** Section in liver of rat from G<sub>4</sub> showing oval cell proliferatin, apoptosis of hepatocytes and mitotic figures (H and E X 200).



**Fig. (10):** Section in liver of rat from G<sub>5</sub> showing no marked histopathological changes (H and E X 200).




**Fig. (11):** Section in liver of rat from G<sub>6</sub> showing kupffer cell proliferation and necrosis of sporadic hepatocytes (H and E X 200).

**Kidneys:**


Microscopically, kidneys of rat from G<sub>1</sub> (vehicle treated control) revealed normal histological structure of renal parenchyma (Fig. 12). Also, kidneys of rat from G<sub>2</sub> which received PP extract revealed apparent normal renal parenchyma (Fig. 13).

Mean while, kidneys of rat from G<sub>3</sub> which treated by low AFS dose showed vacuolations of renal tubular epithelium (Fig. 14) and interstitial nephritis (Fig. 15). Moreover, section from G<sub>4</sub> which treated with high AFS dose revealed tubulointerstitial nephritis (Fig. 16, 17), cystic dilatation of some renal tubules (Fig.16) and dilatation of Bowmen's space (Fig. 17).

However, kidneys of G<sub>5</sub> (treated by low AFS dose and PP extract) revealed no histological changes (Fig. 18). While examined kidneys of rat from G<sub>6</sub> which treated with high AFS dose and PP extract showed vacuolations of renal tubular epithelium (Fig. 19) and focal tubular necrosis associated with leucocytic cells infiltration (Fig. 20).



**Fig. (12):** Section in kidney of rat from G<sub>1</sub> showing the normal histological structure of renal parenchyma (H and E X 200).



**Fig. (13):** Section in kidney of rat from G<sub>2</sub> showing apparent normal renal parenchyma (H and E X 200).




Fig. (14): Section in kidney of rat from G<sub>3</sub> showing vacuolations of renal tubular epithelium (H and E X 200).




Fig. (15): Section in kidney of rat from G<sub>3</sub> showing interstitial nephritis (H and E X 200).




Fig. (16): Section in kidney of rat from G<sub>4</sub> showing interstitial nephritis and cystic dilatation of some renal tubules (H and E X 200)




Fig. (17): Section in kidney of rat From G<sub>4</sub> showing interstitial nephritis and dilatation of Bowman's space (H and E X 200).



Fig. (18): Section in kidney of rat from G<sub>5</sub> showing no histopathological changes (H and E X 200).





Fig. (19): Section in kidney of rat From G<sub>6</sub> showing vacuolations of renal tubular epithelium (H and E X 200).



**Fig. (20):** Section in kidney of rat from G<sub>6</sub> showing focal tubular necrosis associated with leucocytic cells infiltrations (H and E X 200).

## DISCUSSION

Aflatoxin is associated with both toxicity and carcinogenicity in human and animal populations. In developed countries, sufficient amounts of food combined with regulations that monitor aflatoxin levels in these foods protect human populations from significant aflatoxin ingestion. However, in countries where populations are facing starvation or where regulations are either not enforced or nonexistent, routine ingestion of aflatoxin may occur (Cotty *et al.*, 1994 and Abdelhamid, 2010). Worldwide, liver cancer incidence rates are 2 to 10 times higher in developing countries than in developed countries (Henry *et al.*, 1999).

In clinical diagnosis, determination of transaminases is of great importance. The activity of ALT and AST is a sensitive indicator of acute hepatic necrosis and hepatobiliary disease (Kaplan, 1987). In the present study, it was indicated that AFS treatment caused a significant, dose and time-dependent elevation in ALT, AST and ALP activities in serum of rats. The liver is considered to be the principle target organ for AFS and the increase of ALT, AST and ALP activities indicated acute hepatic necrosis (Abdel-Wahhab and Soher, 2003). Similar to our results, Liu *et al.* (2001) reported that administration of rats with AFB<sub>1</sub> by gavage resulting in elevation of serum AST and ALT activities. In addition, Abdel-Wahhab *et al.* (2002) reported that feeding rats with diet contaminated with AFS resulting in significant increase in AST, ALT and ALP in serum. Also Karakilcik *et al.* (2004) fed rabbits on diet AFB<sub>1</sub> for 10 week noticed that, the levels of AST, ALT and ALP in serum were significantly elevated. Moreover, Preetha *et al.* (2006) mentioned that oral administration of rats by AFB<sub>1</sub> resulting in a significant increase in ALP, AST and ALT activity in serum and decreased it in liver after 72h of treatment. The results also showed that the AFS oral administration significantly resulted in decrease of serum level of TP and albumin concentrations with increase of AF dose and time of treatment. These results run in agreement with the data of Dwivedi *et al.* (1993) and Abo-Norag *et al.* (1995) who showed that AF- induced reduction in TP and

albumin in rats. The decrease in plasma protein may be due to degeneration of endoplasmic reticulum and inhibition of protein synthesis (Terao and Ueno, 1978 and Srivastava, 1984). In addition, Osuna and Edds (1982) suggested that such effect may be due to the metabolism of AFS in liver, where it interferes with protein synthesis and RNA production, resulting in decreasing of albumin. Moreover, the low level of TP may indicate protein catabolism and/or kidney dysfunctions (Abdel- Wahhab and Aly, 2003). Data presented herein showed that AFS treatment caused significant elevation of serum creatinine, urea and uric acid in dose and time dependent manner. The present results are supported by the results obtained by Soliman (1994) in rams; El-Zahar et al. (1996) in rabbits and Hassan (1998) in hens. The increased level of blood urea and creatinine with decreased level of blood protein may indicate protein catabolism and /or kidney dysfunction (Abdel-Whhab et al., 1999). Generally, these results indicate degenerative changes and hypofunction of liver and kidneys (Kaplan, 1987).

In the present investigation, treatment of rats by AFS for 6 weeks resulted in hepatotoxicity as revealed by histological study. The degree of severity of pathological lesions depends upon concentration of AFS and time of treatment. Liver of rat treated by low dose of AFS showed vacuolar degeneration of hepatocytes, kupffer cells activation, necrosis of sporadic hepatocytes and focal hepatic necrosis. Meanwhile, liver of rat treated with high AFS dose showed karyomegaly of some nuclei, apoptosis of hepatocytes, portal infiltration with leucocytes, cytokaryomegaly of hepatocytes, oval cells proliferation and mitotic figures. Liver injury by aflatoxicosis was recorded in various studies by Chou et al. (2000); Hiruma et al. (2001) and Preetha et al. (2006) in rats injected by aflatoxin and Hassanein and Abdel Gawad (2001) and Vinita et al. (2003) in rabbits injected by aflatoxin. Kidney of rats was also affected by AFS treatment. The severity of pathological lesions was concentration and time dependent. Kidneys of rats treated with 1mg AFS/ kg b.w. showed vacuulations of renal tubular epithelium and interstitial nephritis. While, kidneys of rats treated with 2mg AFS/ kg b.w. revealed tubulointerstitial nephritis, cystic dilatation of some renal tubules and dilatation of Bowmen's space. Such results were previously recorded by Abdelhamid et al. (1995) in broiler and Vinita et al. (2003) and Abdeen et al. (2004) in rats. Similar alterations were reported by Abdelhamid et al. (2002 a, b and 2005) on rabbits and rats, respectively when fed on aflatoxic diets. Also, Abdelhamid et al. (1990 and 2002 a) came to similar histological changes by aflatoxic rabbits.

The toxic effect produced by AFS on different parameters was explained by Gutteridge and Halliwell (1990) who mentioned that, AFS treatment resulted in enhancement of lipid peroxidation in rats, which is directly related to free radical mediated toxicity. The targets of oxidative damage are usually critical biomolecules such as nucleic acids, proteins, and lipids. In addition, AFS is oxidized by cytochrome P<sub>450</sub> enzymes to AF-8,9oxide (AF- epoxide) which can bind to DNA to form adducts (Choy, 1993). Reaction occurs with high region specificity at the N7 position of guanine residues in DNA (Iyer et al., 1994).

This study has shown that PP extract alone does not have a negative impact on biochemical parameters and histology of liver and kidney.

The current results revealed that co-treatment with PP extract and AFS resulted in a significant improvement in all the tested parameters as indicated by significant alleviation of toxic effect of AFS on ALT, AST, ALP, T.P, albumin, creatinine, urea and uric acid levels in serum. Especially with low AFS dose and at 6<sup>th</sup> week of treatment, but it failed to normalize them except albumin, which restored to normal. The protective effect of PP on liver function against many oxidative stresses was previously recorded by *Toklu et al. (2007)* who reported that PP administration was significantly able to decrease the increase of AST and ALT levels as a result of biliary obstruction induced by bile duct ligation in serum of rats due to its antioxidant and antifibrotic properties. *Parmar and Kar (2007)* mentioned that the treatment of rats with pomegranate peel extract resulted in a decrease of the elevation of serum ALT and creatinine Kinase induced by atherogenic diet. Furthermore, *Masoud and Abo Hagger (2009)* reported that addition of PP powder to diet containing AFS was significantly able to decrease the increase in levels of ALT, AST, ALP, creatinine and urea induced by AFS in serum of guinea pigs.

Present histopathological results revealed that PP extract was able to protect liver and kidney tissue against low AFS dose and minimize the damage of liver and kidney induced by high AFS dose. The beneficial effect of PP extract on tissue of liver and kidneys was previously reported by many studies. *Chidambara Murthy et al. (2002)* reported that feeding rats with methanolic extract of PP extract provides protection against carbon tetrachloride inducing liver injury and restoring the normal hepatic architecture with portal triad, portal veins and hepatic artery and vein were visible. In addition, *Parmar and Kar (2007)* revealed that simultaneous treatment of rats with PP extract ameliorated histological alterations induced by atherogenic diet and liver and kidney tissue were normalized near to those control one. Furthermore, *Toklu et al. (2007)* mentioned that pomegranate peel extract with its antioxidant and antifibrotic properties might be of potential therapeutic value in protecting the liver structure from fibrosis and oxidative injury due to biliary obstruction.

The noticed protective effect of PP extract against toxic effect of AFS could be attributed to its high antioxidants content (*Noda et al., 2002* and *Li et al., 2006*). The phenolic contents of *P. granatum* peel extract acts as a potent free radical scavenger, reducing the levels of hydrogen peroxide and superoxide anion, and consequently inhibit lipid peroxidation, enzyme inactivation, lipid peroxidation of liver and kidney and provide degenerative disease- protective effects on oxidative DNA damage (*Kim et al., 2007* and *Althunibat et al., 2010*). More interesting, *Li et al. (2006)* showed that PP extract had markedly higher antioxidant capacity than pulp extract in scavenging or preventing capacity against superoxide anion, hydroxyl and peroxy radicals as well as inhibiting CuSO<sub>4</sub>-induced LDL oxidation.

In conclusion, the present work suggests that *P. granatum* peel extract might act as a suppressor against tissue damage and inhibit the progression of organs dysfunction induced by AFS.

## REFERENCES

- Abdeen, A.M.; Abdel Hady, E.K.; Mansy, S. E.; Abd-Allah, G.A. and Omar, N. A. (2004): Histopathological and morphological studies on the influence of some aflatoxin on liver, kidneys and testes of albino rat. *Histology and Histochemistry* (44) C: 53-75.
- Abdelhamid, A.M. (2010): Thirty two years (1978-2010) of mycotoxins research at Faculty of Agriculture, Almansoura University, Egypt. *Mycotoxins Technical Articles*, engormix.com, 11p.
- Abdelhamid, A. M.; El-Mansoury, A.M.; Osman, A.I. and El-Azab, S.M. (1999). Mycotoxins as causative for human food poisoning under Egyptian conditions. *J. Agric. Sci., Mansoura Univ.*, 24: 2751-2757.
- Abdelhamid, A. M.; El-Melegy, Kh.M. and Ahmed, A.M. (2005). Possibility of alleviating foodborne aflatoxicosis effects on performance and biochemistry of male Albino white rats. *J. Agric. Sci. Mansoura Univ.*, 30: 822-849.
- Abdelhamid, A.M.; El-Shawaf, I.; Elayoty, S.A.; Ali, M.M. and Gamil, I. (1990). Effect of low level of dietary aflatoxins on baladi rabbits. *Arch. Anim. Nutr., Berlin*, 40:517-537.
- Abdelhamid A.M.; Kandil, W.M.; Arief, H.S.M. and Dorra, T.M. (1995): Effect of some dietary supplements to aflatoxic diets of chicks. III-On the histopathology. *J. Agric. Sci. Mansoura Univ.*, 20:3251-3259.
- Abdelhamid, A.M.; Ragab, M.A. and El-Shaieb, A.F. (2002a). The use of Tafla or aluminosilicate for alleviating toxic effects aflatoxin-contaminated diets of growing rabbits. *Proc, 1<sup>st</sup> Ann. Sc. Conf. Anim. & Fish Prod., Mansoura Fac. Agric.*, 24 & 25 Sep., pp: 389-413.
- Abdelhamid, A.M.; Sallam, A.E; Abd Allah, G.A. and El-Samra, S.H. (2002b). Effect of feeding male rats on aflatoxic diets without or with medicinal herbs (thyme, safflower, ginger, black cumin, and/or garlic). *Proc. 2<sup>nd</sup> Conf. on Food borne Contamination an Egyptian's Health*, April 23-24, Mansoura Fac. Agric., pp: 99-121.
- Abd El-Mageed, F.A. (1987): Some biological and nutritional studies on aflatoxins. M.Sc. Thesis, Zagazig Univ., Fac. of Agric., Egypt.
- Abdel-Wahhab, M.A. and Aly, S.E. (2003): Antioxidants and radical scavenging properties of vegetable extracts in rats fed aflatoxin-contaminated diet. *J. Agric. Food Chem.*, 51 (8): 2409–2414.
- Abdel-Wahhab, M.A.; Nada, S.A. and Amra, H.A. (1999): Effect of aluminosilicate and bentonite on aflatoxin – induced developmental toxicity in rats. *J. Appl. Toxicol.*, 19:199-204.
- Abdel-Wahhab, M.A.; Nada, S.A. and Khalil, F.A. (2002): Physiological and toxicological responses in rats fed aflatoxin-contaminated diet with or without sorbent materials. *Anim. Feed Sci. Technol.*, 97: 209–219.
- Abdel-Wahhab, M.A. and Soher, E.A. (2003): Antioxidants and radical scavenging properties of vegetable extracts in rat feed aflatoxin-contaminated diet. *J. Agric. Food Chem.*, 51: 2409-2414.

- Abo-Norag, M.; Edrington, T.S.; Kubena L.F.; Harvey, R.B. and Phillips T.D. (1995): Influence of a hydrated sodium calcium aluminosilicate and virginiamycin on aflatoxicosis in broiler chickens. *Poult. Sci.*, 74: 626-632.
- Althunibat, O.Y.; Al-Mustafa A.H.; Tarawneh, K.; Khleifat, K.M.; Ridzwan, B.H. and Qaralleh, H. N. (2010): Protective role of *Punica granatum L.* peel extract against oxidative damage in experimental diabetic rats. *Process Bioch.*, 45: 581–585.
- Bennett, J.W. and Klich, M. (2003): Mycotoxins. *Clin. Microbiol., Rev.* 16: 97–516. In: Bintvihok, A., 2002. New insights to controlling mycotoxin anger in ducks. *Feed Tech.*, 6: 28–29.
- Chidambara Murthy, K.N.; Jayaprakasha, G.K. and Singh, R.P. (2002): Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using *in vivo* models. *J. Agric. Food Chem.*, 50: 4791-4791.
- Chou, M.W.; Mikhailova, M.V.; Nichols, J.; Poirier, L.A.; Warbritton, A. and Beland, F.A. (2000): Interactive effect of methyl-deficiency and dietary restriction on liver cell proliferation and telomerase activity in Fischer 344 rats pretreated with aflatoxin B<sub>1</sub>. *Cancer Lett.*, 152 (1): 53-61.
- Choy, W.N. (1993): A review of the dose-response induction of DNA adducts by aflatoxin B<sub>1</sub> and its implications to quantitative cancer – risk assessment. *Mutation Res.*, 296: 181-198
- Cotty, P. J.; Bayman, P.; Egel, D. S. and Elias, K. S. (1994): Agriculture, aflatoxins and *Aspergillus* In: K. A. Powell, A. Renwick, and J. F. Peberdy (ed.). *The genus Aspergillus*. Plenum Press, New York, N.Y., p. 1–27.
- Davis, N.D.; Dioner, V.L. and El-Dridye, D.W. (1966): Production of aflatoxins B<sub>1</sub> and G<sub>1</sub> by *Aspergillus flavus* in a semisynthetic medium. *Appl. Microbiol.*, 14: 378 – 380.
- Do, J.H. and Choi, D.K. (2007): Aflatoxins: Detection, Toxicity, and Biosynthesis. *Biotechnol. Bioprocess. Eng.*, 12: 585-593.
- Dwivedi, Y.; Rastogi R.; Mehrotra R.; Garg, N.K. and Dhawan B.N. (1993): Picroliv protects against aflatoxin B<sub>1</sub> acute hepatotoxicity in rats. *Pharmacol. Res.* 27: 189-199.
- El-Zahar, H.; Tharwat, E.E.; Elaal, W.A.; El-Ashry, M.A.; Saad, M.M. and Amin, S.O. (1996): Rabbit and aflatoxins. 2- Reproductive performance of mature New Zealand white rabbit bucks treated orally with aflatoxins. *Egyptian J. Rabbit Sci.*, 6(1): 67 – 78.
- Gomez, K.A. and Gomez, A.A. (1984): *Statistical Procedure for Agricultural Research*. John Willy and Sons Inc., USA, 96 pp.
- Guo, C. J.; Yang, J. J.; Wei, J. Y.; Li, Y. F.; Xu, J. and Jiang, Y. G. (2003): Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr. Res.*, 23: 1719-1726.
- Gutteridge, J.M.C. and Halliwell, B.(1990): The measurement and mechanism of lipid peroxidation in biological systems. *Trends Biochem. Sci.*, 15:129-135.
- Hassan, O.A. (1998): Effect of aflatoxin B<sub>1</sub>-contaminated diets on some physiological parameters and productive performance of two different layers strain. M.sc. Thesis, Faculty of Agriculture, Alexandria Univeristy, Egypt

- Hassanein, A.M.M. and Abdel Gawad, A.M. (2001): Histological and ultra structural studies on the effect of aflatoxin B<sub>1</sub> on the liver of rabbits. Egypt. J. Zoo., 37: 379-392.
- Henry, S. H.; Bosch, F. X.; Troxell, T. C. and Bolger, P. M. (1999): Reducing liver cancer—global control of aflatoxin. Science, 286:2453–2454.
- Hiruma, S.; Gopalan-Kriczkya, P.; Qina, G.; Gaughanb, J.P. and Lotlikar, P.D. (2001):Differential effects of acetaminophen pretreatment on hepatic aflatoxin B<sub>1</sub>-DNA binding, cellular proliferation, and aflatoxin B<sub>1</sub>-induced hepatic foci in rats and hamsters. Cancer Lett., 170: 117-124.
- Hochstein, P. and Atallah, A. S. (1988): The nature of oxidant and antioxidant systems in inhibition of mutation and cancer. Mutation Res., 202: 363–375.
- Iyer, R.S.; Coles, B.F.; Raney, K.D.; Their, R.; Guengerich, F.P. and Harris, T.M. (1994): DNA adduction by the potent carcinogen aflatoxin B<sub>1</sub>: Mechanistic studies. J. Am. Chem. Soc., 116(5): 1603-1609.
- Julie, J. M.T. (2008): Therapeutic applications of pomegranate (*Punica granatum* L.):A Review. Alternative Medicine Review ,13(2): 128-144.
- Kaplan, M.M. (1987): Laboratory tests. In: Schiff, L., Schiff, E.R. (Eds.), Diseases of the Liver. J. B. Lippincott, Philadelphia, p. 219.
- Karakilcik, A.Z.; Zerim, M.; Arslan, O.; Nazligul, Y and Vural, H. (20004): Effects of vitamin C and E on liver enzymes and biochemical parameters of rabbits exposed to aflatoxin B<sub>1</sub>. Vet. Hum. Toxicol., 46 (4): 190-192.
- Kim, M. R.; Kwon, O.Y. and Sok, D. E. (2007): Antioxidant effects of pomegranate husk (*Granati pericarpium*) in C57BL/6J mice fed high fat and cholesterol diet. FASEB.;21 :844.5
- Lane–Peter, W. and Pearson, A.E.G. (1971): Dietary requirement. Cited in "The Laboratory Animal Principles and Practice". 1<sup>st</sup> Academic press, London and New York. 142.
- Li, Y.; Guo, C.; Yang, J.; Wei, J.; Xu, J. and Cheng, S. (2006): Evaluation of antioxidant properties of pomegranate peel extract in comparison with pomegranate pulp extract. Food Chem., 96: 254-260.
- Liu, J.; Yang, C.F.; Wasser, S.; Shen, H.M.; Looi, C.E. and Ong, C.O. (2001): Protection of *Salvia miltiorrhiza* against aflatoxin-B<sub>1</sub>-induced hepatocarcinogenesis in Fischer 344 rats: Dual mechanisms involved. Life Sci., 69 (3): 309-326.
- Masoud, M.S. and Abo Hagger, Amal A. (2009): Effectiveness of pomegranate peel in amelioration toxic effect of aflatoxin in guinea pigs. J. Agric. Sci. Mansoura Univ., 34(4): 2473-2490.
- Nasr, C. B.; Ayed, N., and Metche, M. (1996): Quantitative determination of the polyphenolic content of pomegranate peel. Zeitschrift für, Lebensmittel Untersuchung und Forschung, 203: 374–378.
- Negi, P.S.; Jayaprakasha, G.K. and Jena, B.S. (2003): Antioxidant and antimutagenic activities of pomegranate peel extracts. Food Chem., 80: 393-397.
- Noda, Y.; Kaneyuki, T.; Mori, A. and Packer, L. (2002): Antioxidant activities of pomegranate fruit extract and its anthocyanidins: dephinidin, cyanidine and pelargonidin. J. Agri. Food Chem., 50:166-171.

- Osuna, O. and Edds, G.T.(1982): Toxicology of aflatoxin B<sub>1</sub>, warfarin, and cadmium in young pigs: clinical chemistry and blood coagulation. *Am. J. Vet. Res.*,43: 387-389.
- Panovska, T. K.; Kulevanova, S. and Stefova, M. (2005): *In vitro* antioxidant activity of some *Teucrium* species (Lamiaceae). *Acta. Pharm.*, 55: 207.
- Parmar, H.S and Kar, A. (2007): Protective role of *Citrus sinensis*, *Musa paradisiaca*, and *Punica granatum* peels against diet-induced atherosclerosis and thyroid dysfunction. *Nutr. Res.*, 27: 710-718.
- Pearse, A.G.W. (1968): Histochemistry, theoretical and applied 3<sup>rd</sup> Ed., Vol. I, Churchill, Livingstone, Edinburgh, London and New York.
- Preetha, S.P.; Kanniappan, M.; Selvakumar, E.; Nagaraj, M. and Varalakshmi, P. (2006): Lupeol ameliorates aflatoxin B<sub>1</sub>-induced peroxidative hepatic damage in rats. *Comp. Biochem. Physiol., Part C Pharmacol. Toxicol.*, 143: 333-339.
- Roos, A. H.; Van der Kamp, H. J. and Marley, E. C. (1997): Comparison of immunoaffinity columns with Florisil/ C18 columns for the determination of aflatoxins in animal feed and maize. *Mycotoxin Research*, 13: 1-10.
- Shotwell, O. L.; Hesseltine, C. W.; Stubblefield, R. D. and Sorenson, W. G. (1966): Production of aflatoxin on rice. *Appl. Microbiol.*, 14: 425-428.
- Soliman, K.M. (1994): Some physiological aspects of aflatoxin treated sheep. Ph. D. Thesis, Faculty of Agriculture, Ain Shams University, Egypt
- Srivastava, A.k.(1984): Pharmacokinetics and therapeutic evaluation of oximes buffalo calves. Ph.D. Thesis, Punjab Agriculture University, Ludhiana, India.
- Tanaka, T.; Kojima, T.; Kawamori, T.; Wang, A.; Suzui, M.; Okamoto, K., and Mori, H. (1993): Inhibition of 4-nitroquinoline-1-oxide-induced rat tongue carcinogenesis by the naturally occurring plant phenolics caffeic, ellagic, chlorogenic and ferulic acids. *Carcinogenesis*, 14: 1321-1325.
- Terao, K. and Ueno, Y. (1978): Morphological and functional damage to cells and tissues. In: *Toxicology, Biochemistry and Pathology of Mycotoxins*, Uraguchi, K. and Adyamazaki, M. (eds.), Kodansha Press, Tokyo, pp. 189-238.
- Toklu, H. Z.; Dumlu, M.U.; Sehirili, O.; Ercan, F.; Gedik, N.; Gokmen, V. and Sener, G.(2007): Pomegranate peel extract prevents liver fibrosis damage in biliary- obstructed rats. *J. Pharmacy and Pharmacol.* 59 (9):1287-1295.
- Ueno, H.; Nakmuro, K.; Sayato, Y. and Okada, S. (1991): Characteristics of mutagenesis by glyoxal in *Salmonella typhimurium*: contribution of singlet oxygen. *Mutation Res.*, 251: 99-107.
- Verma, R. J. (2004): Aflatoxins cause DNA damage. *Int. J. Hum. Genet.* 4(4): 231-236
- Vinita, R.; Prasacl, L.N.; Sinha, B.K. and Rani, V. (2003): Pathological and histochemical changes in liver and kidney in experimental aflatoxicosis in rabbits. *Indian Journal of Veterinary Pathology*, 27: 1-57.
- Waller, R.A. and Duncan, D.B. (1969): A bayes rule for symmetric multiple comparison problem. *Am. State. Assoc. J.*, 64: 1484-1503.

- Wilson, D.M. and Payne, G.A. (1993): Agricultural and veterinary problems: factors affecting *A. flavus* group infection and aflatoxin contamination of crops. In: Eaton, D.L., and Groopman, J.D. (Eds.), The Toxicology of Aflatoxins. Academic Press, New York.
- Yen, G. C. and Chen, H. Y. (1994): Comparison of antimutagenic effect of various tea extracts (green, oolong, pouchong and black tea). J. Food Protection, 57: 54–58.

### تحويل سمية الأفلاتوكسين في الجرذان عن طريق استخدام المستخلص الإيثانولي لقشر الرمان.

أماني عبد المنعم تهامي\* ، أمل عبد العزيز أبو حجر\*\* و ياسر محمد عبد الشفيق\*\*  
\* قسم علم الحيوان و الحشرات-كلية العلوم- جامعة حلوان-حلوان- مصر  
\*\* المركز الإقليمي للأغذية و الأعلاف- مركز البحوث الزراعية – الجيزة-مصر

أجريت الدراسة لمعرفة تأثير المستخلص الإيثانولي لقشر الرمان على خلايا الكبد و الكلى ووظائفهما في الجرذان المعاملة بمستويات مختلفة من الأفلاتوكسين. تم إجراء تجربة لمدة 6 أسابيع على 36 جرذ . تم تقسيم الجرذان الى 6 مجموعات (6جرذان في كل مجموعة). المجموعة الاولى تركت ككنترول (عوملت بالمادة الحاملة) ، المجموعة الثانية عوملت بمستخلص قشر الرمان بالأنبوب المعدى (50مجم / كجم وزن جسم يومياً) ، المجموعة الثالثة عوملت بالأفلاتوكسين عن طريق الأنبوب المعدى (1 مجم / كجم وزن جسم) مرتان في الأسبوع ، المجموعة الرابعة عوملت بالأفلاتوكسين عن طريق الأنبوب المعدى (2 مجم / كجم وزن جسم) مرتان في الأسبوع . المجموعة الخامسة عوملت بالأفلاتوكسين مرتان في الأسبوع (1مجم/ كجم وزن جسم)+ مستخلص قشر الرمان يومياً (50مجم / كجم وزن جسم). المجموعة السادسة عوملت بالأفلاتوكسين مرتان في الأسبوع (2مجم/ كجم وزن جسم)+ مستخلص قشر الرمان يومياً (50مجم / كجم وزن جسم). تم تحليل انزيمات الكبد(ALT, AST, ALP) ، البروتين الكلى و الألبومين ووظائف الكلى (creatinine, urea, uric acid) في المصل عند الأسبوع الثالث و السادس من المعاملة، بينما تم عمل التحليل الباثولوجى للكبد و الكلى في نهاية التجربة . أظهرت النتائج أن كلاً جرعتى الأفلاتوكسين تسببا في زيادة ALT, AST, ALP, creatinine, urea, uric acid و نقص البروتين الكلى و الألبومين بطريقة معتمدة على الجرعة و الوقت. بينما أدت المعاملة بمستخلص قشر الرمان مع الأفلاتوكسين في الحد من التأثير السلبى للأفلاتوكسين خاصة مع الجرعة المنخفضة و الوقت الأطول . أظهرت التحاليل الباثولوجية أن شدة الضرر الحادث في الكبد و الكلى يتناسب طردياً مع جرعات الأفلاتوكسين. بينما كان مستخلص قشر الرمان قادر على حماية الكبد و الكلى من الجرعة المنخفضة من الأفلاتوكسين و تقليل الضرر الحادث للكبد و الكلى بالجرعة المرتفعة من الأفلاتوكسين .

### قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة  
مركز البحوث الزراعية

أ.د / عبد الحميد محمد عبد الحميد  
أ.د / جيهان أحمد حسنى